

Slow mating and equipotential gluing

Wolf Jung, Gesamtschule Aachen-Brand and Jacobs University Bremen.

Joint work with Arnaud Chéritat, Toulouse University.


1. Introduction
2. Equipotential gluing
3. Ideas of the proof of convergence properties
4. Slow mating

Talk at the International Conference on Orthogonal Polynomials and Holomorphic Dynamics, August 16, 2018 in Copenhagen.

1a. Polynomial dynamics

Iteration of $f_c(z) = z^2 + c$. Filled Julia set $\mathcal{K}_c = \{z \in \mathbb{C} \mid f_c^n(z) \not\rightarrow \infty\}$.

External rays \mathcal{R}_c with rational angles land at (pre-)periodic points. The angle is doubled under iteration.





For the Basilica with $f(z) = z^2 - 1$, the relevant angles $1/3$ and $2/3$ are 2-periodic; the rays land at the same fixed point.

For the Rabbit the angles $1/7$, $2/7$, $4/7$ are 3-periodic.

For $f(z) = z^2 + i$ the critical value has the preperiodic angle $1/6$, which is mapped to $1/3$ and $2/3$.

1b. Example of mating

Rational maps or polynomials of higher degree have several critical points. The dynamics is more involved, and the parameter space is higher-dimensional. We may study one-dimensional slices and various combinations.



The first examples of mating were found by Douady–Hubbard. Here a Rabbit and a Basilica are glued along their boundaries, according to conjugate angles.

1c. Definition of matings


Usually assume: $P(z) = z^2 + p$, $Q(z) = z^2 + q$ not in conjugate limbs of \mathcal{M} , with locally connected Julia sets.

Formal mating $g = P \sqcup Q : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ is conjugate to P and Q on half-spheres. Sphere has images of Julia sets and rays, but natural conformal structure only on interior of Julia sets.

Topological mating $P \coprod Q$ shall be a branched cover $S^2 \rightarrow S^2$ obtained from g by collapsing all ray-equivalence classes.

Geometric mating shall be a rational map conjugate to the topological mating, $f \cong P \coprod Q$. The conjugation is conformal on the interior of the filled Julia sets.

1d. Ray-equivalence classes



1e. Construction with Thurston theory

Theorem (Rees–Shishikura–Tan)

Suppose P, Q are postcritically finite (PCF) and not in conjugate limbs (NCL). Then the topological mating $P \amalg Q$ and the geometric mating f exist.

Idea of the construction:

The formal mating $g = P \sqcup Q$ has either no obstructions or only removable obstructions, which surround postcritical ray-equivalence classes.

Modify g to an essential mating \tilde{g} by pinching these ray-equivalence classes. Then \tilde{g} is unobstructed.

If \tilde{g} is not of type $(2, 2, 2, 2)$, the Thurston Theorem provides a combinatorially equivalent rational map f .

The Rees–Shishikura Theorem gives a semi-conjugation Ψ from g to f , which collapses all ray-equivalence classes to distinct points. So f is conjugate to $P \amalg Q$ and the latter exists. ■

1f. Comparison of convergence properties

For a Thurston map g , the pullback gives Homeomorphisms ψ_n and rational maps f_n . In Teichmüller space \mathcal{T} there is a pullback map $\sigma_g : [\psi_n] \mapsto [\psi_{n+1}]$. Note different convergence statements in the case of matings:

For $\sigma_{\tilde{g}}$, we have $[\psi_n] \rightarrow \tau$ and $f_n \rightarrow f$ for general ψ_0 .

For σ_g , we have $\Psi_n \rightarrow \Psi$ for a special choice of Ψ_0 , such that $f_n = f$.

Aim: For σ_g show that $\psi_n \rightarrow \Psi$ and $f_n \rightarrow f$ for more general ψ_0 .

$$\begin{array}{ccc}
 S & \xrightarrow{\psi_0} & S_0 \\
 g \uparrow & & f_0 \uparrow \\
 S & \xrightarrow{\psi_1} & S_1 \\
 g \uparrow & & f_1 \uparrow \\
 S & \xrightarrow{\psi_2} & S_2 \\
 g \uparrow & & f_2 \uparrow \\
 S & \xrightarrow{\psi_3} & S_3 \\
 g \uparrow & & f_3 \uparrow \\
 & \xrightarrow{\dots} &
 \end{array}$$

2a. Equipotential gluing giving spheres

Suppose \mathcal{K}_p and \mathcal{K}_q connected. Equipotential lines are curves with Boettcher coordinates $|\Phi_p(z)| = R$ or $|\Phi_q(z)| = R$, respectively. For $R > 1$, let U'_R and U_R be the domains bounded by equipotential lines of radius R and R^2 , respectively. V'_R and V_R are analogous domains in the dynamic plane of Q .

The annuli in the exterior of the filled Julia sets are mapped to the round annulus $1/R < |z| < R$ by $\Phi_p(z)/R$ and $R/\Phi_q(z)$, respectively. Identify these annuli accordingly to obtain a Riemann surface; it is uniformized to the standard sphere $\mathcal{S}_R = \widehat{\mathbb{C}}$ by $u_R \cup v_R$.

We shall consider images of Julia sets, marked points, external rays of P and Q on \mathcal{S}_R .




Define maps $h_R : \mathcal{S}_e \rightarrow \mathcal{S}_R$ corresponding to $z \mapsto z \cdot |z|^{\log R - 1}$. These are quasiconformal with $K = \log R$ or $K = 1/\log R$. They form a holomorphic motion with respect to $\frac{\log R - 1}{\log R + 1}$ in the unit disk.

2b. Equipotential gluing giving rational maps

The rational map $f_R : \mathcal{S}_{\sqrt{R}} \rightarrow \mathcal{S}_R$ corresponds to the pair of polynomials $P : U_{\sqrt{R}} \rightarrow U_R$ and $Q : V_{\sqrt{R}} \rightarrow V_R$. As $R \rightarrow 1$, it is expected that f_R approximates the geometric mating under suitable conditions.

There is a quasiregular map $g : \mathcal{S}_e \rightarrow \mathcal{S}_e$ such that $f_R \circ h_{\sqrt{R}} = h_R \circ g$.

Now g is conjugate to the formal mating on the filled Julia sets, and equivalent to it in the postcritically finite case. Then $[h_R]$ with $R = R_0^{2^{-t}}$ is a Thurston pullback.



2c. Convergence properties of equipotential gluing

Theorem (Chéritat–J.)

Suppose P and Q are postcritically finite and not from conjugate limbs. Assume $f \cong P \coprod Q$ is not of type $(2, 2, 2)$. Then equipotential gluing with $R \rightarrow 1$ satisfies:

- 1. Marked points converge (with collisions) and $f_R \rightarrow f$.*
- 2. The images of Julia sets, $\mathcal{J}_R = u_R(\partial\mathcal{K}_p) \cup v_R(\partial\mathcal{K}_q)$, satisfy $\mathcal{J}_R \rightarrow \mathcal{J}_f$ with respect to Hausdorff distance.*
- 3. At least if P and Q are hyperbolic, the holomorphic motion converges to the semi-conjugation, $h_R \rightarrow \Psi$. So f is a conformal mating.*

3a. **Convergence of marked points and rational maps**

If g is unobstructed, convergence is a direct application of the Thurston Theorem.

Suppose g has removable obstructions, so the pullback for g diverges in Teichmüller space and moduli space. Then show that marked points collide as expected and converge to common limits. See the general theorem on essential equivalence on page 3b.

It is based on the work of Selinger: he extended σ_g to augmented Teichmüller space $\hat{\mathcal{T}}$, characterized canonical obstructions, and obtained an accumulation statement when the orbifold of a component map is not of type $(2, 2, 2, 2)$. The latter proof employs an interplay of different metrics and the product structure of the canonical boundary stratum.

Use this to construct a suitable path segment in moduli space, which is contained in a neighborhood of a point configuration with repetitions.

Finally, identify eigenvalues of the extended pullback map on configuration space (with additional arguments for identifications with critical points).

3b. General convergence statement

Theorem (J. 2016): *Suppose g is a bicritical Thurston map of degree d , and Γ is a completely invariant multicurve such that:*

- *All components of $S \setminus \Gamma$ except C are disks; the periodic disks are mapped homeomorphically.*
- *C is mapped to itself with degree d ; the essential map \tilde{g} is equivalent to a rational map f .*

Using a normalization at $0, 1, \infty$, such that no disk contains two of them, the Thurston Algorithm for the unmodified map g satisfies:

- *The curves of Γ are pinched, so $[\psi_n]$ diverges in \mathcal{T} .*
- *If f is not of type $(2, 2, 2, 2)$, we have $f_n \rightarrow f$. The marked points converge to (pre-)periodic points of f , identified according to the disks.*

3c. Convergence on Fatou components

Let $R_t = e^{2^{-t}}$ and write f_t for f_{R_t} . Suppose the critical point 0 of P is k -periodic. Then $f_t \circ f_{t+1} \circ \cdots \circ f_{t+k-1}$ has a superattracting fixed point at 0.

Construct a small trapping region around 0, then $\psi_t = h_{R_t}$ converges to a composition of Boettcher conjugations there.

A finite pullback gives locally uniform convergence on every Fatou component of $u_e(\mathcal{K}_p)$, and analogously for Q .

3d. Convergence of Julia sets

Claim: *On any finite set of repelling periodic or preperiodic points, $\psi_t(z)$ converges pointwise.*


Basic idea: use additional obstructions as a tool to prove convergence of additional marked points, cf. page 3e.

To apply the general convergence result from page 3b, we need to show that the component map \widehat{g} is unobstructed and equivalent to f (with additional marked points). Either extend Tan Lei's techniques, or use the semi-conjugation Ψ from g to f according to the Rees–Shishikura Theorem.

To show that \mathcal{J}_f is in an ε -neighborhood of \mathcal{J}_t , cover it with a finite collection of $\varepsilon/2$ -neighborhoods of repelling periodic points, and employ the claim above.

To show that \mathcal{J}_t is in an ε -neighborhood of \mathcal{J}_f , use convergence on Fatou components, together with the fact that there are only finitely many components of diameter $\geq \varepsilon$.

3e. Additional marked points and obstructions



3f. Convergence of holomorphic motions

Here suppose in addition that both P and Q are periodic, so that the orbifold metric ρ_f is expanding for the perturbed maps f_R as well when $R \rightarrow 1$. (For a smaller expansion constant $1 < \lambda' < \lambda$. Here a tiny neighborhood of the critical points, well within the trapping region, is excluded.)

Now consider the ρ_f -length of segments $\{h_{R_t}(z) \mid T \leq t \leq T+1\}$ for z outside of the trapping region. It shrinks exponentially with T , uniformly in z . (Note that the estimate requires two terms, one involving the expanding property of the metric, and one coming from f_t depending on t .)

This gives uniform convergence of h_{R_t} as $t \rightarrow \infty$, or $R \rightarrow 1$, with respect to the ρ_f -metric and the spherical metric. The limit is a semi-conjugation from g to f , which coincides with Ψ from the Rees–Shishikura Theorem.

4a. **Slow mating**

For postcritically finite polynomials P and Q with critical orbits (p_i) and (q_i) , the unmodified Thurston Algorithm for the formal mating $g = P \sqcup Q$ can be implemented with a path in moduli space as follows: Fix $R_1 \geq 5$ and interpolate the radius as $\log(R_t) = 2^{1-t} \log(R_1)$ for $0 \leq t \leq 1$. Set

$$x_i(t) = \frac{1 + (1-t)q/R_1^2}{1 + (1-t)p/R_1^2} \cdot \frac{p_i/R_t}{1 + (1-t)q/R_1^4(p_i - p)} \approx \frac{p_i}{R_t} \quad \text{and}$$

$$y_i(t) = \frac{1 + (1-t)q/R_1^2}{1 + (1-t)p/R_1^2} \cdot \frac{R_t \left(1 + (1-t)p/R_1^4(q_i - q) \right)}{q_i} \approx \frac{R_t}{q_i} .$$

The initial path for $0 \leq t \leq 1$ can be pulled back uniquely for $1 < t < \infty$, choosing the sign below by continuity.

$$z_i(t+1) = \pm \sqrt{\frac{1 - y_1(t)}{1 - x_1(t)} \cdot \frac{z_{i+1}(t) - x_1(t)}{z_{i+1}(t) - y_1(t)}} \quad \text{for } t \geq 0 .$$

4b. Slow mating approximates equipotential gluing

Suppose \mathcal{K}_p and \mathcal{K}_q are connected. So p and q may be postcritically infinite, in conjugate limbs, non-locally connected, or giving type $(2, 2, 2, 2)$.

Proposition: *As $R \rightarrow \infty$, we have $u_R(z) \sim z/R$ and $v_R(z) \sim R/z$ on U'_R and V'_R , respectively.*

Proof by a composition of quasi-conformal maps with small dilatation.

Corollary: *For all $t \geq 0$, compare slow mating at time t to equipotential gluing at radius $R = R_0^{2^{-t}}$. Then h_R is uniformly close to slow mating in the sense of \mathcal{T} , $\widehat{\mathcal{T}}$, for a suitably initialized ψ_t , or considering marked points in $\widehat{\mathbb{C}}^{|Z|-3}$.*

In the postcritically infinite case, slow mating makes limited sense here as an approximation for a finite time with a large number of marked points.