

+

+

Constitution and Stability of Atoms and Molecules

J.P. Solovej
Aarhus University

Ascona July 1996

+

1

+

+

OVERVIEW OF LECTURES

Schrödinger Theory (QM)

- Definition of ground state energy
 - ionization energy
 - binding energy
 - stability
 - radius of atoms
 - size of molecules
- Review of known (rigorous) results
- Qualitative features of the periodic table and their possible formulations as conjectures

Hartree-Fock Theory (HF)

- Differences and similarities with QM
- Validity as approximation

Thomas-Fermi Theory (TF)

- No-binding
- Sommerfeld asymptotics
- Validity as approximation (semiclassics)
- Consequences for QM and HF

+

2

+

+

I. The Schrödinger Theory (QM) of Atoms and Molecules:

Molecule of N electrons and K nuclei. Nuclear charges $\underline{Z} = (Z_1, \dots, Z_K)$. Hamiltonian:

$$H_{N,\underline{Z}}^{\text{tot}} = H = \sum_{k=1}^K T_k + \sum_{i=1}^N t_i + V_C.$$

The nuclear kinetic energy operator:

$$T_k = -\frac{1}{2M_k} \Delta_{R_k}, \quad M_k = \text{Nuclear mass}$$

The electron kinetic energy operator:

$$t_i = -\frac{1}{2} \Delta_{x_i}.$$

The Coulomb potential V_C is the function

$$V_C = - \sum_{k=1}^K \sum_{i=1}^N Z_k |x_i - R_k|^{-1} + \sum_{i < j} |x_i - x_j|^{-1} \\ + \sum_{k < \ell} Z_k Z_\ell |R_k - R_\ell|^{-1}$$

Units: $m_e = \hbar = e$.

+

3

+

+

The Pauli principle: The electrons are Fermions.
The physical Hilbert space is

$$\mathcal{H}_{\text{physical}}^{\text{tot}} = \bigotimes_k^K L^2(\mathbb{R}_{R_k}^3) \otimes \bigwedge_i^N L^2(\mathbb{R}^3; \mathbb{C}^2).$$

$H_{N,\underline{Z}}^{\text{tot}}$ is Friderichs' extended from C_0^∞ functions.

Removing the Center of Mass Motion:

Define $H_{N,\underline{Z}}$:

$$H_{N,\underline{Z}}^{\text{tot}} = -(2(N + \sum_k M_k))^{-1} \Delta_{\text{CM}} \otimes \mathbf{1} + \mathbf{1} \otimes H_{N,\underline{Z}}$$

$H_{N,\underline{Z}}$ acts in the space $\mathcal{H}_{\text{physical}}$ such that

$$\mathcal{H}_{\text{physical}}^{\text{tot}} = L^2(\mathbb{R}_{\text{CM}}^3) \otimes \mathcal{H}_{\text{physical}}.$$

$\mathcal{H}_{\text{physical}}$ is a subspace (because of the symmetry properties) of the Hilbert space of \mathbb{C}^{2N} -valued L^2 functions on a $\mathbb{R}^{3(N+K)-1}$ dimensional space.

Definition: Ground State Energy:

$$E^{\text{QM}}(N, \underline{Z}) := \inf \text{Spec}_{\mathcal{H}_{\text{physical}}} H_{N,\underline{Z}}$$

+

4

+

+

Definition: Atomic and Molecular Stability:

$E^{\text{QM}}(N, \underline{Z})$ is an eigenvalue of $H_{N, \underline{Z}}$.

Strong Stability: $E^{\text{QM}}(N, \underline{Z})$ is a discreet eigenvalue.

HVZ Criterion: Strong Stability follows if

$$E^{\text{QM}}(N, \underline{Z}) < \inf \{ E^{\text{QM}}(N', \underline{Z}') + E^{\text{QM}}(N'', \underline{Z}'') : \\ N' + N'' = N, \underline{Z}' \oplus \underline{Z}'' = \underline{Z} \}$$

“It costs energy to break into two clusters.”

Result: The neutral molecule H_2 (i.e., $(N, Z_1, Z_2) = (2, 1, 1)$) is stable.

Refr: Richard, Fröhlich, Graf & Seifert, *PRL* '93.

Bounds on Stability for diatomic molecules:
 N cannot be too small. Refr: Ruskai, *LMP* '89
Sol., *CMP* '90

+

+

+

Definition: Stability of Matter:

$$E^{\text{QM}}(N, \underline{Z}) \geq C(\max Z_k)(N + K)$$

Refr: Dyson&Lenard, *JMP*'67&'68,
Lieb&Thirring, *PRL*'75.

Infinite Nuclear Mass Approximation:

$$H(N, \underline{Z}, \underline{R}) := \sum_{i=1}^N t_i + V_c$$

$\underline{R} = (R_1, \dots, R_K)$ parameters. $H(N, \underline{Z}, \underline{R})$ acts
in $\mathcal{H}(N) = \bigwedge^N L^2(\mathbb{R}^3; \mathbb{C}^2)$.

Born-Oppenheimer Energy Function:

$$E^{\text{QM}}(N, \underline{Z}, \underline{R}) := \inf \text{spec}_{\mathcal{H}(N)} H(N, \underline{Z}, \underline{R})$$

Ground State Energy:

$$E^{\text{QM}}(N, \underline{Z}) := \inf_{\underline{R}} E(N, \underline{Z}, \underline{R})$$

Stability: There exists \underline{R}_0 such that

- (i) $E^{\text{QM}}(N, \underline{Z}, \underline{R}_0) = E^{\text{QM}}(N, \underline{Z})$
- (ii) $E^{\text{QM}}(N, \underline{Z}, \underline{R}_0)$ is an eigenvalue of $H(N, \underline{Z}, \underline{R})$.
(Notice translation invariance.)

+

+

+

Ground State Configuration: (R_0, ψ) with $\psi \in \mathcal{H}(N)$ ground state eigenfunction for $H(N, \underline{Z}, \underline{R}_0)$.

Density:

$$\rho(x) = N \int \cdots \int \|\psi(x, x_2, \dots, x_N)\|_{\mathbb{C}^{2N}}^2 dx_2 \cdots dx_N$$

Binding energy:

$$BE^{\text{QM}}(N, \underline{Z}) :=$$

$$\inf\{E^{\text{QM}}(N', \underline{Z}') + E^{\text{QM}}(N'', \underline{Z}'')\} - E^{\text{QM}}(N, \underline{Z})$$

Infimum over $N' + N'' = N$ and $\underline{Z}' \oplus \underline{Z}'' = \underline{Z}$ where both \underline{Z}' and \underline{Z}'' are not empty.

Ionization Energy:

$$IE^{\text{QM}}(N, \underline{Z}) = E^{\text{QM}}(N - 1, \underline{Z}) - E^{\text{QM}}(N, \underline{Z}).$$

OBS: Both BE and IE are non-negative.

Strong Stability: (i) $BE^{\text{QM}}(N, \underline{Z}) > 0$

(ii) $IE^{\text{QM}}(N, \underline{Z}) > 0$.

HVZ criterion: Strong Stability \Rightarrow Stability.

+

7

+

+

- Result: Atoms with $N < Z$ are stable.
- Result: H^- (i.e. $(N, Z) = (2, 1)$) is stable. (variational proof).
- Result: H^{--} (i.e., $(N, Z) = (3, 1)$) is unstable

The last result which was an open problem for many years is a corollary of

Theorem 1 (Lieb, *PRL*'84 & *PRA*'84):
If a molecule with (N, \underline{Z}) is stable then

$$N < 2 \sum_{k=1}^K Z_k + K.$$

The proof for atoms ($K=1$) is remarkably simple. It relies on Hardy's inequality:

$$-\operatorname{Re} \int \bar{f} |x| \Delta f = \int |\nabla |x|^{1/2} f(x)|^2 dx - \frac{1}{4} \int |x|^{-2} \left| |x|^{1/2} f(x) \right|^2 dx \geq 0$$

and a very clever use of the triangle inequality.

+

8

+

+

Result: Neutral molecules are *always* stable (in the infinite mass approximation). This is a corollary of

Theorem 2 (van der Waals attraction)

Lieb&Thirring *PRA*'86

If $N = \sum_k Z_k$, $\underline{Z} = \underline{Z}' \oplus \underline{Z}''$, $N' = \sum_k Z'_k$, and $N'' = \sum_k Z''_k$ and $\max_{k,\ell} |R_k - R_\ell|$ large enough

$$E^{\text{QM}}(N, \underline{Z}, \underline{R}) \leq E^{\text{QM}}(N', \underline{Z}') + E^{\text{QM}}(N'', \underline{Z}'') - C(\underline{Z}', \underline{Z}'') \left(\max_{k,\ell} |R_k - R_\ell| \right)^{-6}.$$

Radius of Atoms: Define \mathfrak{R}_ν by

$$\int_{|x-R_1| \geq \mathfrak{R}_\nu} \rho(x) dx = \nu.$$

The radius is naturally defined as \mathfrak{R}_1 , the radius to last electron.

Size of Molecules: $D = \max_{k,\ell} |R_k - R_\ell|$.

Important Conjecture For Atoms: The ionization energy $IE(N, Z)$ is non-increasing in N .

+

+

+

Hartree-Fock (HF) Theory: Consider only

$$\Psi = \varphi_1 \wedge \cdots \wedge \varphi_N, \quad \varphi_i \in L^2(\mathbb{R}^3; \mathbb{C}^2). \quad (1)$$

Hartree-Fock energy function:

$$E^{\text{HF}}(N, \underline{Z}, \underline{R}) = \inf_{\Psi \text{ as in (1)}} \frac{(\Psi, H(N, \underline{Z}, \underline{R})\Psi)}{(\Psi, \Psi)}$$

Reformulation: $\gamma = \text{proj. on span}\{\varphi_1, \dots, \varphi_N\}$.
 2×2 -matrix Kernel $\gamma(x, y)$.

$$\frac{(\Psi, H(N, \underline{Z}, \underline{R})\Psi)}{(\Psi, \Psi)} = \mathcal{E}_{\underline{Z}, \underline{R}}^{\text{HF}}(\gamma) = \text{Tr} \left[\left(-\frac{\Delta}{2} - \sum_{k=1}^K \frac{Z_k}{|x - R_k|} \right) \gamma \right] \\ + \mathcal{D}(\gamma) - \mathcal{E}\mathcal{X}(\gamma) + \sum_{k < \ell} \frac{Z_k Z_\ell}{|R_k - R_\ell|}$$

Direct energy:

$$\mathcal{D}(\gamma) = \frac{1}{2} \iint \rho_\gamma(x) |x - y|^{-1} \rho_\gamma(y) dx dy.$$

Density:

$$\rho(x) = \rho_\gamma(x) = \text{Tr}_{\mathbb{C}^2}[\gamma(x, x)].$$

Exchange energy:

$$\mathcal{E}\mathcal{X}(\gamma) = \frac{1}{2} \iint \frac{\text{Tr}_{\mathbb{C}^2}[|\gamma(x, y)|^2]}{|x - y|} dx dy.$$

+

+

+

Lieb's variational principle (PRL'81):

$$E^{\text{HF}}(N, \underline{Z}, \underline{R}) = \inf \{ \mathcal{E}_{\underline{Z}, \underline{R}}^{\text{HF}}(\gamma) : 0 \leq \gamma \leq 1, \text{Tr} \gamma = N \}$$

We now proceed as in QM:

Ground state energy:

$$E^{\text{HF}}(N, \underline{Z}) = \inf_{\underline{R}} E^{\text{HF}}(N, \underline{Z}, \underline{R})$$

Stability: *There exists \underline{R}_0 such that*

$$(i) E^{\text{HF}}(N, \underline{Z}, \underline{R}_0) = E^{\text{HF}}(N, \underline{Z})$$

(ii) *There exists γ with $\text{Tr} \gamma = N$ such that*

$$E^{\text{HF}}(N, \underline{Z}, \underline{R}_0) = \mathcal{E}_{\underline{Z}, \underline{R}_0}^{\text{HF}}(\gamma).$$

Note that eigenvalue (for QM) has been replaced by existence of minimizer.

Just as for QM we have the quantities:

$$BE^{\text{HF}}(N, \underline{Z}), IE^{\text{HF}}(N, \underline{Z}), \rho^{\text{HF}}(x), \mathfrak{R}_\nu^{\text{HF}}$$

(Almost) all previous (infinite nuclear mass) results hold.

Atomic stability for $N < Z + 1$: Lieb & Simon CMP '77.

Exception: Van der Waals Theorem.

+

+

+

Remark: As for QM the HF minimizer need not be unique: For atoms the spherical symmetry may be broken: atoms may have angular momentum and spin.

The mean-field operator: For γ with $0 \leq \gamma \leq 1$

$$H_{\gamma}^{MF} = -\frac{1}{2}\Delta - \sum_{k=1}^K Z_k |x - R_k|^{-1} \\ + \rho_{\gamma} * |x|^{-1} - \mathcal{K}_{\gamma}$$

$$\rho_{\gamma}(x) = \text{Tr}_{\mathbb{C}^2}[\gamma(x, x)], \quad \mathcal{K}_{\gamma}(x, y) = \frac{\gamma(x, y)}{|x - y|}$$

Theorem 3 (Self-consistency:) *If γ minimizes $\mathcal{E}_{\underline{Z}, \underline{R}}^{\text{HF}}$ with $\text{Tr}\gamma = N$ then γ is the projection onto the "lowest" eigenfunctions of H_{γ}^{MF} .*

Theorem 4 (No unfilled shells) *(Bach, Lieb, Loss & Sol. PRL '94): If γ is a HF minimizer then the space of the N "lowest" eigenfunctions of H_{γ}^{MF} is unique !*

+

12

+

+

When is HF a good approximation to QM?
Morally: When the density is big!

Theorem 5 *If $N \rightarrow \infty$, $\|\underline{Z}\| \rightarrow \infty$, $N/\|\underline{Z}\|$ fixed then (clearly $E^{\text{HF}}(N, \underline{Z}) \rightarrow -\infty$) $\exists \varepsilon > 0$ s.t.*

$$\left| \frac{E^{\text{QM}}(N, \underline{Z})}{E^{\text{HF}}(N, \underline{Z})} - 1 \right| \leq c |E^{\text{HF}}(N, \underline{Z})|^{-\frac{2}{7} - \varepsilon}.$$

References:

For atoms: Fefferman, Seco Bull AMS '90

For molecules: Bach CMP '93. Improved in Graf, Sol. Rev. Math. Phys. '94

If $D(H, \underline{Z})$ and $Ex(N, \underline{Z})$ denote the direct and exchange energies then in the limit above

$$\frac{D(N, \underline{Z})}{E^{\text{HF}}(N, \underline{Z})} = o(1),$$

$$\frac{Ex(N, \underline{Z})}{E^{\text{HF}}(N, \underline{Z})} = o(|E^{\text{HF}}(N, \underline{Z})|^{-\frac{2}{7}})$$

+

+

+

Thomas-Fermi Theory:

$$\mathcal{E}_{\underline{Z}, \underline{R}}^{\text{TF}}(\rho) = \frac{3(3\pi^2)^{2/3}}{10} \int \rho^{5/3} - \sum_{k=1}^K Z_k \int |x - R_k|^{-1} \rho(x) dx$$

$$+ \frac{1}{2} \iint \rho(x) |x - y|^{-1} \rho(y) dx dy + \sum_{1 \leq k < l \leq K} Z_k Z_l |R_k - R_l|^{-1}$$

Energy Function:

$$E^{\text{TF}}(N, \underline{Z}, \underline{R}) = \inf \{ \mathcal{E}_{\underline{Z}, \underline{R}}^{\text{TF}}(\rho) \mid \rho, \int \rho = N \}.$$

The next theorem from Brezis&Lieb CMP'79 states that Molecules are never stable in TF theory:

Theorem 6 (strong no-binding) *Given N, \underline{Z} and \underline{R} there exists a decomp. $N = N_1 + \dots + N_K$ such that*

$$E^{\text{TF}}(N, \underline{Z}, \underline{R}) \geq \sum_{k=1}^K E^{\text{TF}}(N_k, Z_k, R_k) + C \min \{ \|\underline{Z}\|^{7/3}, |\underline{R}_k - R_l|^{-7} \}$$

+

+

+

Only neutral and subneutral atoms are stable:

Theorem 7 (Lieb&Simon Adv.Math.'77)

For $K = 1$ there exists a minimizer ρ^{TF} for $\mathcal{E}_{Z,R}^{\text{TF}}$ with $\int \rho^{\text{TF}} = N$ if and only if $N \leq Z$.

Effective Potential for atoms:

$$\varphi^{\text{TF}}(x) = Z|x|^{-1} - \rho^{\text{TF}} * |x|^{-1}, \quad R = 0$$

TF equation: φ^{TF} is the unique positive solution to

$$\Delta \varphi^{\text{TF}} = 2^{7/3}(3\pi)^{-1} \varphi_{\text{TF}}^{3/2} - 4\pi Z \delta_0$$

Consequence:

Theorem 8 (Sommerfield Asymptotics)

$$0 \leq 3^4 2^{-3} \pi^2 |x|^{-4} - \varphi_{\text{TF}}(x) \leq CZ^{(4-\tau)/3} |x|^{-\tau},$$

where $\tau = (1 + \sqrt{73})/2 \approx 4.8$, $4 - \tau \approx -0.8$.

The universal infinite TF atom:

$$\varphi_{\text{TF}}(x) \rightarrow 3^4 \cdot 2^{-3} \pi^2 |x|^{-4} \text{ as } Z \rightarrow \infty.$$

+

+

+

For large Z , TF theory is a good approximation to HF theory and then in turn to QM.

Theorem 9 (Lieb&Simon Adv.Math.'77)

If $N/\|\underline{Z}\|$ fixed $\exists \varepsilon > 0$ such that

$$\left| \frac{E^{\text{HF}}(N, \underline{Z}, \underline{R})}{E^{\text{TF}}(N, \underline{Z}, \underline{R})} - 1 \right| \leq C |E^{\text{TF}}(N, \underline{Z}, \underline{R})|^{-\varepsilon}$$

Same with HF replaced by QM.

This is a semiclassical approximation:

$$\text{Tr}_{L^2(\mathbb{R}^3; \mathbb{C}^2)} \left[\left(-\frac{1}{2} h^2 \Delta - V \right)_- \right] = \frac{2^{5/2}}{15\pi^2} h^{-3} \int V^{5/2} + o(h^{-3}), V \geq 0$$

The Legendre transform of

$$V \mapsto \frac{2^{5/2}}{15\pi^2} V^{5/2}$$

is

$$\rho \mapsto \frac{3}{10} (3\pi^2)^{2/3} \rho^{5/3}$$

which appeared in the TF functional.

+

+

+

Consequences for QM of no-binding and neutrality:

- Lieb-Thirring proof of stability of matter
- Atoms are asymptotically neutral:

Theorem 10 : *If $Z_n \rightarrow \infty$, atom (N_n, Z_n) stable then*

$$\limsup_{n \rightarrow \infty} \frac{N_n}{Z_n} \leq 1$$

References:

Lieb&Sigal&Simon&Thirring CMP'88

Fefferman&Seco CMP'90

Seco&Sigal&Solovej CMP'90

Compare: *Without the Pauli principle there exists $0 < \gamma < 1$ such that if $N(Z)$ denotes the maximal number of electrons such that (N, Z) is stable then $\lim \frac{N(Z)}{Z} = 1 + \gamma$.*

" \geq " Benguria&Lieb PRL'83,

" \leq " Solovej LMP'90

+

+

+

Consequence for QM of strong no-binding (Brezis-Lieb law): Molecules are asymptotically neutral.

Theorem 11 : If $\underline{Z}_n = Z_n \underline{z}$ with $\sum \underline{z} = 1$, $Z_n \rightarrow \infty$, molecule (N_n, \underline{Z}_n) stable then

$$\lim \frac{N_n}{Z_n} = 1.$$

(Note upper *and* lower bound).

References:

Solovej CMP'90. (diatomic case)

Ruskai-Solovej, Springer Lecture Notes in Physics **403** (polyatomic case).

Bounds on the size of molecules are also derived.

+

+

+

Consequence for HF of Sommerfeld asymptotics: in the HF theory the radius of atoms is bounded above and below independently of Z

Theorem 12 (Solovej '96) : *Consider the HF effective potential*

$$\varphi^{\text{HF}}(x) = Z|x|^{-1} - \rho^{\text{HF}} * |x|^{-1}, R_1 = 0,$$

$$\rho^{\text{HF}}(x) = T_{R_{\mathbb{C}^2}}[\gamma^{\text{HF}}(x, x)].$$

There exist universal constants, $\varepsilon, D > 0$ such that

$$0 < |x| \leq D \Rightarrow |\varphi^{\text{HF}}(x) - \varphi^{\text{TF}}(x)| \leq |x|^{-4+\varepsilon},$$

+

+

+

Open problems

- Monotonicity of ionization energies
- Understand the effect of relativistic corrections
- Prove bounds uniform in Z on the radius of QM atoms and the size of QM molecules
- Likewise bounds on the binding energies and ionization energies
- Prove bounds on binding energies and size of molecules in HF (still open)
- Prove upper bounds uniform in Z on the density of matter both in HF and QM.

+