

UNIVERSAL OPERATIONS IN HOCHSCHILD HOMOLOGY

NATHALIE WAHL

ABSTRACT. We provide a general method for finding all natural operations on the Hochschild complex of \mathcal{E} -algebras, where \mathcal{E} is any algebraic structure encoded in a prop with multiplication, as for example the prop of Frobenius, commutative or \mathcal{A}_∞ -algebras. We show that the chain complex of all such natural operations is approximated by a certain chain complex of *formal operations*, for which we provide an explicit model that we can calculate in a number of cases. When \mathcal{E} encodes the structure of open topological conformal field theories, we identify this last chain complex, up to quasi-isomorphism, with the moduli space of Riemann surfaces with boundaries, thus establishing that the operations constructed by Costello and Kontsevich-Soibelman via different methods identify with all formal operations. When \mathcal{E} encodes open topological quantum field theories (or symmetric Frobenius algebras) our chain complex identifies with Sullivan diagrams, thus showing that operations constructed by Tradler-Zeinalian, again by different methods, account for all formal operations. As an illustration of the last result we exhibit two infinite families of non-trivial operations and use these to produce non-trivial higher string topology operations, which had so far been elusive.

INTRODUCTION

To any dg-algebra, or more generally \mathcal{A}_∞ -algebra A , one can associate the Hochschild chain complex $C_*(A, A)$ of the algebra with coefficients in itself (as in e.g. [21, 7.2.4]). Many authors have constructed operations on this complex, i.e. natural maps

$$\nu_A : C_*(A, A)^{\otimes p} \rightarrow C_*(A, A)^{\otimes q}$$

for some $p, q \geq 0$, under various assumptions on the type of A , see e.g. [24, Chap 4],[29] or [21].

In the present paper, we systematically address the question of finding the chain complex $\text{Nat}_\mathcal{E}^\otimes(p, q)$ of *all* such natural operations, for various classes of \mathcal{A}_∞ -algebras with extra structure \mathcal{E} (associative, commutative, Frobenius, ...). We do this by introducing a chain complex $\text{Nat}_\mathcal{E}(p, q)$ of all *formal operations*, loosely speaking obtained by forgetting a symmetric monoidal condition in the definition of the algebras, and provide an explicit model for the chain complex of all formal operations (Theorem A). We are able to identify this chain complex up to quasi-isomorphism in terms of well-known objects, in several interesting cases (see Theorems B and C). The chain complex of formal operations comes with a chain map $r : \text{Nat}_\mathcal{E}(p, q) \rightarrow \text{Nat}_\mathcal{E}^\otimes(p, q)$ to the natural operations. While this map is not an isomorphism in general, we do show that it has good properties, under various assumptions on \mathcal{E} , e.g., if \mathcal{E} comes from an operad. And for a general \mathcal{E} we identify the natural operations in terms of the formal operations of the completion $\widehat{\mathcal{E}}$ of \mathcal{E} via an isomorphism $\text{Nat}_\mathcal{E}^\otimes(p, q) = \text{Nat}_{\widehat{\mathcal{E}}}^\otimes(p, q) \xleftarrow{\cong} \text{Nat}_{\widehat{\mathcal{E}}}(p, q)$. Before detailing our computations, we start by making precise which types of algebraic structures we consider.

An \mathcal{A}_∞ -algebra, as originally defined by Stasheff [33], can be described as an enriched symmetric monoidal functor $\Phi : \mathcal{A}_\infty \rightarrow \text{Ch}$ from a certain dg-category \mathcal{A}_∞ with objects the natural numbers, to Ch , the category of chain complexes over \mathbb{Z} : the \mathcal{A}_∞ -algebra

associated to such a functor Φ is its value at 1, together with multiplications m_i , $i \geq 2$, encoded as morphisms in the category \mathcal{A}_∞ .

Let \mathcal{E} be any symmetric monoidal dg-category with objects the natural numbers (i.e. a prop, also denoted PROP [26, §24]), equipped with a symmetric monoidal dg-functor $i : \mathcal{A}_\infty \rightarrow \mathcal{E}$ which is the identity on objects—we call such a pair (\mathcal{E}, i) a *prop with \mathcal{A}_∞ -multiplication*. An \mathcal{E} -algebra is defined to be a strong symmetric monoidal functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$, i.e. a functor equipped with natural isomorphisms $\Phi(1)^{\otimes n} \xrightarrow{\cong} \Phi(n)$ compatible with the symmetries of \mathcal{E} . If (\mathcal{E}, i) is a prop with \mathcal{A}_∞ -multiplication, i endows $\Phi(1)$ with the structure of an \mathcal{A}_∞ -algebra. We define the Hochschild complex of the \mathcal{E} -algebra Φ with respect to i as the standard Hochschild complex of the \mathcal{A}_∞ -algebra $\Phi(1)$, i.e.

$$\mathbf{C}_*(\Phi) := C_*(\Phi(1), \Phi(1)) = \bigoplus_{n \geq 1} \Phi(1)^{\otimes n}$$

with the standard Hochschild differential defined using the multiplication and higher multiplications of $\Phi(1)$.

A natural operation, with p inputs and q outputs, on the Hochschild complex of \mathcal{E} -algebras is the data of a linear map

$$\nu_\Phi : \mathbf{C}_*(\Phi)^{\otimes p} \rightarrow \mathbf{C}_*(\Phi)^{\otimes q}$$

for any \mathcal{E} -algebra Φ , which is natural with respect to maps of \mathcal{E} -algebras. The set of such linear maps between chain complexes is again canonically a chain complex. In other words, if we let $\text{Fun}^\otimes(\mathcal{E}, \text{Ch})$ denote the category of \mathcal{E} -algebras and $C_\mathcal{E}^{\otimes p} : \text{Fun}^\otimes(\mathcal{E}, \text{Ch}) \rightarrow \text{Ch}$ the functor taking an \mathcal{E} -algebra Φ to the p th power of its Hochschild complex $\mathbf{C}_*(\Phi)^{\otimes p}$, the natural operations, with p inputs and q outputs, on the Hochschild complex of \mathcal{E} -algebras is the chain complex $\text{Hom}(C_\mathcal{E}^{\otimes p}, C_\mathcal{E}^{\otimes q})$.

More generally, we want to consider operations of the form

$$\nu_\Phi : \mathbf{C}_*(\Phi)^{\otimes n_1} \otimes \Phi(1)^{\otimes m_1} \rightarrow \mathbf{C}_*(\Phi)^{\otimes n_2} \otimes \Phi(1)^{\otimes m_2}$$

allowing also copies of the algebra $\Phi(1)$ itself as input and output. Generalising the previous paragraph, the chain complex of those $(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$ -operations is hence given by

$$\text{Nat}_\mathcal{E}^\otimes(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}) := \text{Hom}(C_\mathcal{E}^{\otimes(n_1, m_1)}, C_\mathcal{E}^{\otimes(n_2, m_2)})$$

where $C_\mathcal{E}^{\otimes(n, m)} : \text{Fun}^\otimes(\mathcal{E}, \text{Ch}) \rightarrow \text{Ch}$ is the functor taking Φ to $\mathbf{C}_*(\Phi)^{\otimes n} \otimes \Phi(1)^{\otimes m}$.

One can in fact define a Hochschild complex $\mathbf{C}_*(\Phi)$ for *any* functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$ (not necessarily strong symmetric monoidal) by setting

$$\mathbf{C}_*(\Phi) := \bigoplus_{n \geq 1} \Phi(n),$$

noting that the usual Hochschild differential is still well-defined. Letting

$C_\mathcal{E}^{(n, m)} : \text{Fun}(\mathcal{E}, \text{Ch}) \rightarrow \text{Ch}$ more generally denote the analogous extension of $C_\mathcal{E}^{\otimes(n, m)}$, with value on Φ defined as a sum of terms $\Phi(k_1 + \cdots + k_n + m)$, we can hence consider the chain complex of all the natural $(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$ -operations

$$\text{Nat}_\mathcal{E}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}) := \text{Hom}(C_\mathcal{E}^{(n_1, m_1)}, C_\mathcal{E}^{(n_2, m_2)})$$

on the Hochschild complex of generalized \mathcal{E} -algebras.

There is a restriction map

$$r : \text{Nat}_\mathcal{E}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}) \longrightarrow \text{Nat}_\mathcal{E}^\otimes(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$$

which in general need neither be injective, surjective, nor a quasi-isomorphism. We think of the left-hand side as *formal operations* on \mathcal{E} -algebras, and refer to them as such. We show that the prop \mathcal{E} can always be replaced by a new prop $\widehat{\mathcal{E}}$, its *completion*, with same category of algebras and satisfying

$$\text{Nat}_\mathcal{E}^\otimes(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}) \cong \text{Nat}_{\widehat{\mathcal{E}}}^\otimes(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}) \cong \text{Nat}_{\widehat{\mathcal{E}}}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix}),$$

so natural operations can always be described as formal operations for a different prop (see Definition 2.5 and Corollary 2.10)—the completion $\widehat{\mathcal{E}}$ is in fact the prop whose morphisms are $\widehat{\mathcal{E}}(m_1, m_2) = \text{Nat}_{\widehat{\mathcal{E}}}^{\otimes}([\![_{m_1}^0], [\![_{m_2}^0]$).

We show that the restriction map r is injective (resp. surjective or a quasi-isomorphism) for all $([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]])$ if and only if it is injective (resp. surjective or a quasi-isomorphism) for all $([\![_{m_1}^0], [\![_{m_2}^0])$; or, said differently, it is injective if and only if any non-zero morphism $f \in \mathcal{E}(m_1, m_2)$ acts non-trivially on some \mathcal{E} -algebra, it is surjective if any operation on \mathcal{E} -algebras is induced by \mathcal{E} , and it is a quasi-isomorphism if and only if $\mathcal{E} \rightarrow \widehat{\mathcal{E}}$ is a quasi-isomorphism (see Theorem 2.9 for the first two statements and Corollary 2.2 for the last one). The map r is for example always injective if \mathcal{E} is the prop associated to an operad, and one can show that it is an isomorphism when restricting to operads and algebras in vector spaces, over a field of characteristic 0 [10] (see Examples 2.11 and 2.13 for more details).

Our main technical theorem, Theorem 2.1, gives an explicit description of the chain complex $\text{Nat}_{\mathcal{E}}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]])$:

Theorem A (see Theorem 2.1). *For any prop with \mathcal{A}_{∞} -multiplication (\mathcal{E}, i) , there is an isomorphism of chain complexes*

$$\text{Nat}_{\mathcal{E}}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]]) \cong \prod_{j_1, \dots, j_{n_1} \geq 1} \bigoplus_{k_1, \dots, k_{n_2} \geq 1} \mathcal{E}(j + m_1, k + m_2)[k - j + n_1 - n_2],$$

where $j = j_1 + \dots + j_{n_1}$, $k = k_1 + \dots + k_{n_2}$, and where the differential on the right hand side is the sum of the differential of \mathcal{E} , a multi-Hochschild and multi-coHochschild differential. The square brackets indicate a shift in grading.

Theorem 2.1 gives in addition a version for the reduced Hochschild complex when \mathcal{E} has units. A version for higher Hochschild homology (in the sense of [32]) can be found in [18].

Using a spectral sequence for the right hand side, a corollary of the theorem is that $\text{Nat}_{\mathcal{E}}$ is homotopy invariant in \mathcal{E} (see Corollary 2.2).

We use the same spectral sequence to identify the homology of $\text{Nat}_{\mathcal{E}}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]])$ in three cases detailed below: the case of unital \mathcal{A}_{∞} -algebras, the case of Frobenius algebras, and the case of open field theories. Klamt computed in addition the case of commutative algebras in [17], and, to a large extent, the case of commutative Frobenius algebras in [19]. To further exemplify our approach, we show in Proposition 2.14 how the cap product in Hochschild homology can be seen as part of the chain complex of formal operations $\text{Nat}_{\mathcal{E}}([\![_0^1], [\![_0^1])$ for $\mathcal{E} = \text{End}(A)$ the endomorphism prop of the algebra A considered.

Now, consider the case where $\mathcal{E} = \mathcal{O}$, the open cobordism category. It is a prop with morphisms $\mathcal{O}(m_1, m_2)$ a chain model given in terms of fat graphs for the moduli space of Riemann cobordisms from m_1 closed unit intervals to m_2 closed unit intervals. There is a natural inclusion $i : \mathcal{A}_{\infty} \rightarrow \mathcal{O}$ induced by including forests into all graphs, so \mathcal{O} is a prop with \mathcal{A}_{∞} -multiplication. Furthermore, let $\mathcal{OC}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]])$ denote a chain model for the moduli space of Riemann cobordisms from n_1 circles and m_1 intervals to n_2 circles and m_2 intervals, where each component of the cobordism has at least one incoming or one free (i.e. neither incoming nor outgoing) boundary component; see Section 3.1 for precise definitions. Theorem 3.1 provides the following description of the formal operations on the Hochschild complex of open field theories:

Theorem B (see Theorem 3.1). *There is an inclusion*

$$\mathcal{OC}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]]) \xrightarrow{\sim} \text{Nat}_{\mathcal{O}}([\![_{m_1}^{n_1}], [\![_{m_2}^{n_2}]])$$

which is a split-injective quasi-isomorphism.

That the open-closed cobordism category \mathcal{OC} can be used to construct natural operations on the Hochschild complex of \mathcal{O} -algebras was first discovered by Costello and Kontsevich–Soibelman via a priori different constructions [4, 21]. What we show here is that these, up to quasi-isomorphism, both give all *formal* operations. This gives an answer to a wish expressed in [4, 1.3] of a natural algebraic characterization of the category of chains on moduli spaces of curves in terms of the functors which assign to an \mathcal{O} -algebra the tensor powers of its Hochschild chains.

We also obtain a computation of the formal operations in the case of unital \mathcal{A}_∞ -algebras, i.e. $\mathcal{E} = \mathcal{A}_\infty^+$, from the computation of \mathcal{O} , using that the prop of unital \mathcal{A}_∞ -algebras \mathcal{A}_∞^+ is a subcategory of \mathcal{O} . In fact $\text{Nat}_{\mathcal{A}_\infty^+}([m_1]^{n_1}, [m_2]^{n_2})$ is a union of components of $\text{Nat}_{\mathcal{O}}([m_1]^{n_1}, [m_2]^{n_2})$, namely those associated to cobordisms which are a union of certain annuli and discs, generated up to the above quasi-isomorphism by the well-known operations, namely the \mathcal{A}_∞ -structure of the algebra, the inclusion of the algebra in its Hochschild complex, and Connes–Rinehart’s boundary operator B (see Theorem 3.7).

Finally we consider the case $\mathcal{E} = H_0(\mathcal{O})$, the 0th homology of \mathcal{O} . Symmetric monoidal functors $H_0(\mathcal{O}) \rightarrow \text{Ch}$ are also known as open topological quantum field theories, which by [23, Cor 4.5] correspond precisely to symmetric Frobenius algebras. In [35], Tradler and Zeinalian show that a certain chain complex of Sullivan diagrams acts on the Hochschild cochain complex of symmetric Frobenius algebras. (See also [36] where the dual action on the Hochschild chains is constructed.) Our Theorem 3.8 shows that these define all formal operations in the following sense:

Theorem C (see Theorem 3.8). *There is an inclusion*

$$\mathcal{SD}([m_1]^{n_1}, [m_2]^{n_2}) \xrightarrow{\sim} \text{Nat}_{H_0(\mathcal{O})}([m_1]^{n_1}, [m_2]^{n_2})$$

which is a split-injective quasi-isomorphism where $\mathcal{SD}([m_1]^{n_1}, [m_2]^{n_2})$ is a chain complex of Sullivan diagrams on n_2 circles, with n_1 “incoming” boundary cycles and $m_1 + m_2$ labeled leaves.

The chain complex $\mathcal{SD}([m_1]^{n_1}, [m_2]^{n_2})$ is a quotient of $\mathcal{OC}([m_1]^{n_1}, [m_2]^{n_2})$ with the same H_0 , and is briefly recalled in Section 3.3. It computes the homology of the harmonic compactification of moduli space [8].

Beyond providing a model for the right-hand side, Theorem C can also be used “in reverse” to give new information about the left-hand side: The theorem indeed implies that the Hochschild homology of Frobenius algebras can be used as representation of the homology of Sullivan diagrams, and in Section 4 we use this representation for the Frobenius algebra $H^*(S^n)$ to produce two infinite families of non-trivial homology classes in Sullivan diagrams of increasing degree (non-trivial both with integral and rational coefficient). The above cycles furthermore give rise to rational higher string operations, since for any 1-connected manifold M there is an $H_0(\mathcal{O}, \mathbb{Q})$ -algebra $A(M)$ whose Hochschild homology is isomorphic to $H^*(LM, \mathbb{Q})$ (using [22] and [16], see also [9] or [36, Sec 6.6]), and these operations are non-trivial since they are non-trivial on $HH_*(H^*(S^n, \mathbb{Q}), H^*(S^n, \mathbb{Q})) \cong H^*(LS^n, \mathbb{Q})$ by the same computation. As far as we know, these are the first non-trivial higher string topology operations constructed. (Note also that our higher degree cycles live in components of arbitrarily high genus and number of incoming boundary components, which should be contrasted to a result of Tamanai [34] that states that for large genus or number of incoming boundaries, the degree 0 string topology operation on $H^*(LM)$ constructed in [3, 12] are zero.)

The paper is organised as follows: A technical reformulation of Theorem A is the statement that formal operations on the Hochschild complex of \mathcal{E} -algebras are given by taking an iterated Hochschild and then coHochschild construction on the category \mathcal{E} . Section 1 defines the Hochschild and coHochschild complex, and their reduced versions, in the generality needed in this paper, and establishes basic properties of these, which

are used in Section 2 to give a proof of Theorem A. In Section 2 we also compare the formal operations to the natural operations, and give the cap-product example mentioned earlier. In Section 3 we identify the formal operations in the case of open field theories, Frobenius algebras, and \mathcal{A}_∞ -algebras, proving in particular Theorems B and C, building on our joint work with Westerland [36]. The main ingredient in the proof in each case is a computation of the homology of the relevant iterated coHochschild constructions in terms of chain complexes associated to certain cosimplicial sets of partitions. In the process we need the general fact that the homology of the canonical chain complex associated to a cosimplicial set is concentrated in degree 0, the proof of which we postpone to Section 5. Finally Section 4 gives examples of non-trivial operations on the Hochschild complex of Frobenius algebras, and in particular of non-trivial higher string operations. Section 0 sets up the notations and conventions about functors and dg-categories used in the paper, and the final Section 6 recalls how to define complexes of fat graphs, and exemplifies their use in the context of \mathcal{A}_∞ -algebras.

Acknowledgment. This paper is the continuation of the author's joint work with Craig Westerland [36]. In fact, this paper was started as an elaboration on an suggestion of Craig that the coHochschild complex could play a role. Many other people have influenced this paper with comments, questions and answers. In particular, I would like to thank Alexander Berglund, Bill Dwyer, Benoît Fresse, Anssi Lahtinen and Bruno Vallette. I would furthermore like to thank the referee for many helpful suggestions. The author was supported by the Danish National Sciences Research Council (DNSRC) and the European Research Council (ERC), as well as by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92). The author would also like to thank IHES for its hospitality during the final stages of the redaction of this article.

CONTENTS

Introduction	1
0. Functors and dg-categories; notations and conventions	5
1. The Hochschild and coHochschild complexes	6
1.1. Homotopy invariance	10
1.2. Reduced complexes	11
2. Formal and natural operations on the Hochschild complex	14
2.1. Formal operations	14
2.2. Restriction to natural operations	17
2.3. Example: the cap product	20
3. Three computations of the formal operations	22
3.1. Open topological conformal field theories	22
3.2. Unital \mathcal{A}_∞ -algebras	28
3.3. Strict Frobenius algebras	29
4. Examples of non-trivial higher string topology operations	31
5. Appendix A: The chain complex associated to a cosimplicial set	33
6. Appendix B: Graph complexes	35
References	36

0. FUNCTORS AND DG-CATEGORIES; NOTATIONS AND CONVENTIONS

We work throughout this paper with dg-categories, that is categories enriched over chain complexes, by which we mean differential graded \mathbb{Z} -modules. These are categories \mathcal{C} whose morphism sets $\mathcal{C}(n, m)$ are chain complexes and such that composition in the

category is given by chain maps

$$\mathcal{C}(n, p) \otimes \mathcal{C}(p, m) \longrightarrow \mathcal{C}(n, m).$$

We will denote by Ch the dg-category of chain complexes, i.e. the category whose objects are chain complexes and whose morphisms are linear maps of any degree. We denote by $\text{Hom}(V, W)$ the chain complex of morphisms from V to W in Ch . The differential on $\text{Hom}(V, W)$ is defined so that the evaluation map $V \otimes \text{Hom}(V, W) \rightarrow W$ is a chain map for every pair of chain complexes V, W . Explicitly, for $f \in \text{Hom}(V, W)$, this gives $df(v) = (-1)^{|v|}(d_W(f(v)) - f(d_V(v)))$. In particular, chain maps $V \rightarrow W$ are the degree 0 cycles in $\text{Hom}(V, W)$.

A dg-functor $F : \mathcal{C} \rightarrow \text{Ch}$ is an enriched functor, that is a functor such that the maps

$$F(n) \otimes \mathcal{C}(n, m) \longrightarrow F(m)$$

are chain maps.

Note that our convention in this paper is that morphisms act on the right. This has an influence on the signs we work with. If a reader wants to compare our signs to those one would obtain having the morphisms act on the left, this can be done by multiplying with the Koszul signs coming from permuting the factors appropriately before and, if relevant, after the map or identification considered.

Let \mathcal{C} be a small dg-category. Given dg-functors $F, G : \mathcal{C} \rightarrow \text{Ch}$, let

$$\text{Hom}_{\mathcal{C}}(F, G) \subset \prod_{k \in \text{Obj}(\mathcal{C})} \text{Hom}(F(k), G(k))$$

denote the chain complex of natural transformations $F \rightarrow G$.

Given dg-functors $F : \mathcal{C} \rightarrow \text{Ch}$ and $G : \mathcal{C}^{op} \rightarrow \text{Ch}$, we denote by

$$F \otimes_{\mathcal{C}} G = \bigoplus_{k \in \text{Obj}(\mathcal{C})} F(k) \otimes G(k) / \sim$$

the tensor product of F and G , where the equivalence relation is given by $F(f)(x) \otimes y \sim (-1)^{|y||f|} x \otimes G(f)(y)$ for any $x \in F(k)$, $y \in G(l)$ and $f \in \mathcal{C}(k, l)$. This is a chain complex with differential $d = d_F + d_G$ (with the usual Koszul sign convention).

In this paper, functors describing algebras will be monoidal functors. Note that those will always be *strong monoidal* (also called split monoidal), i.e. satisfying that $F(n) \otimes F(m) \rightarrow F(n + m)$ is an isomorphism.

By a *quasi-isomorphism of functors* with values in chain complexes, we mean a natural transformation given by pointwise quasi-isomorphisms.

1. THE HOCHSCHILD AND COHOCHSCHILD COMPLEXES

In this section, we recall from [36] our generalization of the Hochschild complex of a dg-algebra, and define its dual, a generalization of the coHochschild complex of a dg-coalgebra. We then show that our coHochschild complex is homotopy invariant, and show the equivalence between the reduced and unreduced (co)Hochschild constructions when the algebras have units. The homotopy invariance of the Hochschild complex is proved in [36, Prop. 5.7].

We recall briefly the prop \mathcal{A}_{∞} , already mentioned in the introduction. We will use the language of graphs recalled in the Appendix Section 6. \mathcal{A}_{∞} is a dg-category with objects the natural numbers $\mathbb{N} = \{0, 1, 2, \dots\}$ and with morphisms $\mathcal{A}_{\infty}(n, m)$ the free \mathbb{Z} -module on the set of graphs which are unions of m planar trees, with a total of $n = n_1 + \dots + n_m$ *incoming* labeled leaves, with each $n_i \geq 1$, in addition to the root of each tree, considered here as an outgoing leaf. (See Figure 1(a) for an example.) From $n = 0$, there is only the identity morphism. As detailed in Section 6, $\mathcal{A}_{\infty}(n, m)$ can be given the structure of a chain complex using the valence minus 3 of vertices to define

the degree, and the sum of all possible blow-ups to define the differential. Composition in \mathcal{A}_∞ is defined by gluing outgoing leaves to incoming leaves with the same label, and disjoint union of trees defines a symmetric monoidal structure on \mathcal{A}_∞ .

Recall that an \mathcal{A}_∞ -algebra is a dg-module A equipped with multiplications m_2, m_3, \dots where m_k is a degree $k-2$ map $m_k : A^{\otimes k} \rightarrow A$. These maps satisfy relations among them which make m_2 into a multiplication associative up to all higher homotopies. The category defined above is called \mathcal{A}_∞ because symmetric monoidal functors $\Phi : \mathcal{A}_\infty \rightarrow \text{Ch}$ correspond exactly to (non-unital) \mathcal{A}_∞ -algebras. Such a functor Φ will satisfy $\Phi(n) = A^{\otimes n}$ for some dg-module A , and the multiplication m_k is obtained by evaluating Φ on the morphism in $\mathcal{A}_\infty(k, 1)$ defined by a tree with a single vertex (see Figure 1(b)). The relations satisfied by the m_k 's follow from the dg-structure of \mathcal{A}_∞ and the fact that Φ is a dg-functor. (See Example 6.1 and e.g. [36, 3.1] and [25, C.2 and 9.2.7] for more details.)



FIGURE 1. Morphism in $\mathcal{A}_\infty(6, 2)$ and the map $m_k \in \mathcal{A}_\infty(k, 1)$.

Dually, an \mathcal{A}_∞ -coalgebra is a functor $\Psi : \mathcal{A}_\infty^{op} \rightarrow \text{Ch}$.

We will define the (co)Hochschild complex of a (co)algebra using a functor

$$\mathcal{L} : \mathcal{A}_\infty^{op} \longrightarrow \text{Ch}$$

which we now define. As a graded module, $\mathcal{L}(m) = \bigoplus_{n \geq 1} \mathcal{A}_\infty(m, n)[n-1]$, where $[n-1]$ indicates a shift in degree by $[n-1]$, and \mathcal{A}_∞ acts on \mathcal{L} by precomposition. (Here we use the notation $V[n]$ for the graded module defined by $V[n]_* = V_{*-n}$.) To describe the differential, we write $\mathcal{L}(m)$ as

$$\mathcal{L}(m) = \bigoplus_{n \geq 1} \mathcal{A}_\infty(m, n) \otimes L_n$$

where $L_n = \langle l_n \rangle$ the free module on a single generator l_n in degree $n-1$. The differential on $\mathcal{L}(m)$ is the sum of the differential $d_{\mathcal{A}_\infty}$ of the first factor with a twisted differential d_L coming from the second factor, which we describe now, first pictorially and then algebraically.

The pictorial way to define the differential is to think of the generator l_n of L_n as an oriented fat graph with a single (white) vertex and n ordered leaves attached to it. The differential of l_n is then the sum of all ways of blowing up that vertex, just like we blew up trees in \mathcal{A}_∞ , with the only difference that we blow up a white vertex as a pair of a white and a black vertex, and that the white vertex is allowed to have valence 1 or 2. Figure 2 shows the graph l_3 and all its blow-ups. As a blow-up of l_n will always be of




FIGURE 2. Differential applied to l_3 .

the form a tree attached to an l_k with $1 \leq k < n$, we can write the differential as a map

$$d_L : L_n \rightarrow \bigoplus_{1 \leq k < n} \mathcal{A}_\infty(n, k) \otimes L_k.$$

This then induces a self-map of $\mathcal{L}(m)$ using composition in \mathcal{A}_∞ :

$$d_L : \mathcal{L}(m) = \bigoplus_{n \geq 1} \mathcal{A}_\infty(m, n) \otimes L_n \rightarrow \bigoplus_{n, k \geq 1} \mathcal{A}_\infty(m, n) \otimes \mathcal{A}_\infty(n, k) \otimes L_k \rightarrow \bigoplus_{k \geq 1} \mathcal{A}_\infty(m, k) \otimes L_k$$

We can describe d_L explicitly in terms the \mathcal{A}_∞ -multiplications m_k . We give here the description without signs. The signs are more easily thought of as orientations of graphs. (See Example 6.1 for how to interpret graph orientations as signs.) We can decompose the differential as

$$d_L(l_n) = \sum_{k < n} f_{n,k} \otimes l_k$$

where $f_{n,k} \in \mathcal{A}_\infty(n, k)$ is itself given as

$$(1) \quad f_{n,k} = \sum_{i=1}^n m_{n,k}^i = \sum_{i=1}^n \pm m_{n-k+1}^i$$

with $m_{n-k+1}^i \in \mathcal{A}_\infty(n, k)$ the morphism that multiplies with m_{n-k+1} the entries $i, i+1, \dots, i+n-k$ (considered modulo n) and $m_{n,k}^i = \pm m_{n-k-1}^i$ defined to contain the sign/orientation it receives in the differential. The term $m_{n,k}^i$ corresponds, in the pictorial description, to the graph obtained from l_n by blowing up its leaved labeled $i, i+1, \dots, i+n-k$ (modulo n) to form a tree with one vertex attached to l_k at the i th position.

Remark 1.1. We can now think of an element of $\mathcal{L}(m)$ as a sum of graphs with m leaves obtained by attaching trees (the elements of $\mathcal{A}_\infty(m, n)$) to a white vertex (l_n), as in Figure 3. Such graphs are particular examples of black and white graphs in the language of [36] and the differential of $\mathcal{L}(m)$ is the differential of black and white graphs as defined in Section 2 of that paper. (See also Section 3.1.)




FIGURE 3. Element in $\mathcal{L}(12)$, decomposable as an element of $\mathcal{A}_\infty(12, 5) \otimes L_5$.

We are now ready to define our version of the Hochschild complex. Let \mathcal{E} be a monoidal dg-category. Given a dg-functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$ and an object $m \in \mathcal{E}$, we can define a new functor

$$\Phi(- + m) : \mathcal{E} \rightarrow \text{Ch}$$

by setting $\Phi(- + m)(n) = \Phi(n + m)$ and $\Phi(- + m)(f) = \Phi(f + id_m)$. Note that for any morphism $g \in \mathcal{E}(m, m')$, $\Phi(id + g)$ induces a natural transformation $\Phi(- + m) \rightarrow \Phi(- + m')$.

Recall that we call a pair $\mathcal{E} = (\mathcal{E}, i)$ a prop with \mathcal{A}_∞ -multiplication if \mathcal{E} is a symmetric monoidal dg-category and $i : \mathcal{A}_\infty \rightarrow \mathcal{E}$ is a symmetric monoidal dg-functor which is the identity on objects.

Definition 1.2. Let (\mathcal{E}, i) be a prop with \mathcal{A}_∞ -multiplication. For a dg-functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$, define its Hochschild complex as the dg-functor $C(\Phi) : \mathcal{E} \rightarrow \text{Ch}$ given on objects by

$$C(\Phi)(m) := i^* \Phi(- + m) \otimes_{\mathcal{A}_\infty} \mathcal{L}$$

and on morphisms by

$$C(\Phi)(f) := i^* \Phi(id + f) \otimes_{\mathcal{A}_\infty} id.$$

Dually, for a dg-functor $\Psi : \mathcal{E}^{op} \rightarrow \text{Ch}$, we define its coHochschild complex as the dg-functor $D(\Psi) : \mathcal{E}^{op} \rightarrow \text{Ch}$ given on objects by

$$D(\Psi)(m) := \text{Hom}_{\mathcal{A}_\infty^{op}}(\mathcal{L}, (i^{op})^* \Psi(- + m))$$

and on morphisms by

$$D(\Psi)(f)(\underline{h}) := (i^{op})^* \Psi(id + f) \circ \underline{h}.$$

The Hochschild and coHochschild complexes are natural in Φ and Ψ and hence define self-maps of the functor categories:

$$C : \text{Fun}(\mathcal{E}, \text{Ch}) \rightarrow \text{Fun}(\mathcal{E}, \text{Ch}) \quad \text{and} \quad D : \text{Fun}(\mathcal{E}^{op}, \text{Ch}) \rightarrow \text{Fun}(\mathcal{E}^{op}, \text{Ch}).$$

We denote by $C^p = C \circ \dots \circ C$ and $D^p = D \circ \dots \circ D$ the iterated functors.

We can describe $C(\Phi)$ and $D(\Psi)$ more explicitly as follows. Since \mathcal{L} is quasi-free,

$$\begin{aligned} C(\Phi)(m) &\cong \bigoplus_{n \geq 1} \Phi(n+m) \otimes L_n &\cong \bigoplus_{n \geq 1} \Phi(n+m)[n-1] \\ D(\Psi)(m) &\cong \prod_{n \geq 1} \text{Hom}(L_n, \Psi(n+m)) &\cong \prod_{n \geq 1} \Psi(n+m)[1-n] \end{aligned}$$

as a graded modules, where the second isomorphism in each case comes from the fact that each L_n is generated by a single element in degree $n-1$. The differential in the first case is given, for $x \in \Phi(n+m)$, by

$$d(x \otimes l_n) = d_\Phi x \otimes l_n + (-1)^{|x|} \sum_{k=1}^{n-1} \Phi(i(f_{n,k}) + id_m)(x) \otimes l_k$$

with $f_{n,k}$, $k < n$, the terms of the differential of L_n as in equation (1) above. In the second case, an element \underline{h} of degree d in $D(\Psi)(m)$ is determined by a sequence $(\underline{h}(l_1), \underline{h}(l_2), \dots) \in \prod_{n \geq 1} \Psi(n+m)_{n-1+d}$. In this notation, the differential is given by

$$d\underline{h}(l_n) = (-1)^{n-1} \left(d_\Psi(\underline{h}(l_n)) - \sum_{k=1}^{n-1} \Psi(i(f_{n,k}) + id_m)(\underline{h}(l_k)) \right).$$

Remark 1.3. For $\Phi : \mathcal{A}_\infty \rightarrow \text{Ch}$ strong symmetric monoidal, with $\Phi(n) = A^{\otimes n}$ for A a dg- (\mathcal{A}_∞) -algebra, the value of $C(\Phi)$ at 0 is the Hochschild complex of A :

$$C(\Phi)(0) = \bigoplus_{n \geq 1} \Phi(n) = \bigoplus_{n \geq 1} A^{\otimes n}$$

with differential the sum of the differential of A and the Hochschild differential which multiplies, with higher and higher multiplications, all possible pairs, triples, quadruples, ..., of consecutive factors thought of as cyclically arranged on a circle (see [36, Rem 5.2]).

Similarly, when $\Psi : \mathcal{A}_\infty^{op} \rightarrow \text{Ch}$ is the functor associated to a coalgebra C concentrated in degree 0, i.e. $\Psi(n) = C^{\otimes n}$ with \mathcal{A}_∞^{op} -structure given by the comultiplication of C , the chain complex $D(\Psi)(0)$ recovers the classical coHochschild complex of C . Indeed, we have

$$D_*(\Psi)(0) \cong \prod_{n \geq 1} C^{\otimes n}[-n+1].$$

If C is concentrated in degree 0, then $D(\Psi)(0)$ is only non-zero in non-positive degrees, with elements of degree $-d$ of the form $(0, \dots, 0, a_0 \otimes \dots \otimes a_d, 0, \dots)$. The differential $d = d_L$ takes such an element to the element of degree $-d-1$ whose only non-zero term is

$$(-1)^d \sum_{i=0}^d (-1)^i a_0 \otimes \dots \otimes (\sum a'_i \otimes a''_i) \otimes \dots \otimes a_d - \sum a''_0 \otimes a_1 \otimes \dots \otimes a_d \otimes a'_0$$

where $\sum a'_i \otimes a''_i$ is the comultiplication of a_i (see for example [6, Sec 3.1]).

Example 1.4. Suppose A is an \mathcal{A}_∞ -algebra, and let $\Phi : \mathcal{A}_\infty \rightarrow \text{Ch}$ with $\Phi(n) = A^{\otimes n}$ be the functor defining this \mathcal{A}_∞ -structure. Consider the functor $\text{Hom}_A : \mathcal{A}_\infty^{op} \rightarrow \text{Ch}$ defined by

$$\text{Hom}_A(n) = \text{Hom}(A^{\otimes n}, A)$$

and, for $f \in \mathcal{A}_\infty(m, n)$, by $\text{Hom}_A(f) : \text{Hom}(A^{\otimes n}, A) \rightarrow \text{Hom}(A^{\otimes m}, A)$ induced by precomposition with $\Phi(f)$. Then the coHochschild complex of Hom_A evaluated at 0 is

$$D(\text{Hom}_A)(0) = \prod_{n \geq 1} \text{Hom}(A^{\otimes n}, A)[1 - n]$$

though with a differential which is *not* that of the Hochschild cochains—the coHochschild differential in $D(\text{Hom}_A)$ cyclically precomposes with multiplications. In Section 2.3, we will though give an embedding of the Hochschild cochain complex $C^*(A, A)$ inside the complex $D(C(\text{Hom}_A)(0))(0)$.

For symmetric monoidal functors, the iterated Hochschild complex computes tensor powers of the Hochschild complex of the associated algebra (see [36, Prop. 5.10]). For the coHochschild complex, as a graded vector space,

$$D^n(\Phi)(m) = \prod_{k_1, \dots, k_n \geq 1} \text{Hom}(L_{k_1} \otimes \dots \otimes L_{k_n}, \Phi(k_1 + \dots + k_n + m))[n - k_1 - \dots - k_n].$$

The functors $C^n(-)(m)$ and $D^n(-)(m)$ can be constructed as homology theories associated to a union of n circles and m points in the same way classical Hochschild homology is associated to the circle (in Pirashvili's much more general language of higher Hochschild homology [32]).

1.1. Homotopy invariance. Proposition 5.6 in [36] shows that the Hochschild complex functor is homotopy invariant. We show in this section the less straightforward fact that the coHochschild complex also has this property.

Let $\Psi : \mathcal{E}^{op} \rightarrow \text{Ch}$ be a dg-functor. Each chain complex $D(\Psi)(m)$ admits a natural filtration: using the identification $D(\Psi)(m) \cong \prod_{q \geq 1} \Psi(q + m)[1 - q]$, we define a filtration by

$$F^s = \prod_{q \geq s} \Psi(q + m)[1 - q].$$

Each F^s is indeed a subcomplex as the differential of $D(\Psi)$ is of the form $d = d_\Psi + d^L$, with d_Ψ the differential of Ψ not affecting q , and d^L increasing q . We have

$$D := D(\Psi)(m) = F^1 \supset F^2 \supset \dots \supset F^s \supset F^{s+1} \supset \dots$$

is an exhaustive filtration ($D = \bigcup_s F^s$), which is moreover complete ($\lim_s D/F^s = \prod_s \Psi(s + m) = D$). We have $F^s/F^{s+1} \cong \Psi(s + m)[1 - s]$

Note that the coHochschild part of the differential $d^L = \sum_{n > q} f_{n,q}$ takes F^s to F^{s+1} , so that the spectral sequence associated to this filtration has the form

$$E_{-p,q}^1 = H_q(\Psi(p + 1 + m), d_\Psi) \implies H_{q-p}(D(\Psi)(m)).$$

Proposition 1.5. *Let (\mathcal{E}, i) be a prop with \mathcal{A}_∞ -multiplication and let $\Psi, \Psi' \in \text{Fun}(\mathcal{E}^{op}, \text{Ch})$. A quasi-isomorphism $\eta : \Psi \xrightarrow{\cong} \Psi'$ induces a quasi-isomorphism $D(\eta) : D(\Psi) \xrightarrow{\cong} D(\Psi')$.*

Recall that by a quasi-isomorphism of functors with values in chain complexes, we mean a natural transformation given by pointwise quasi-isomorphisms.

Proof. A natural transformation $\Psi \rightarrow \Psi'$ induces a natural transformation $D(\Psi) \rightarrow D(\Psi')$ and we are left to show that this natural transformation is by quasi-isomorphisms. For each m , we use the filtration of $D(\Psi)(m)$ and $D(\Psi')(m)$ defined above, and the associated spectral sequence. A quasi-isomorphism of functors $\Psi \xrightarrow{\cong} \Psi'$ induces an

isomorphism of the E^1 -terms of the spectral sequences. The result then follows from the Eilenberg-Moore Comparison Theorem [37, Thm 5.5.11]. \square

1.2. Reduced complexes. Let \mathcal{A}_∞^+ be the dg-category obtained from \mathcal{A}_∞ by adding one generating morphism $u \in \mathcal{A}_\infty^+(0, 1)$ with the relations that it acts as a unit for m_2 , i.e. we have $m_2 \circ (u + id) = id = m_2 \circ (id + u) \in \mathcal{A}_\infty^+(1, 1)$, and such that $m_k(id_i + u + id_{k-i-1}) = 0$ for all $k > 2$ and $0 \leq i \leq k - 1$. Then symmetric monoidal dg-functors $\Phi : \mathcal{A}_\infty^+ \rightarrow \text{Ch}$ corresponds exactly to \mathcal{A}_∞ -algebras $A = \Phi(1)$ equipped with a strict unit for the multiplication $m_2 : A \otimes A \rightarrow A$.

If the functor $i : \mathcal{A}_\infty \rightarrow \mathcal{E}$ extends to a functor $i^+ : \mathcal{A}_\infty^+ \rightarrow \mathcal{E}$ from the prop of unital \mathcal{A}_∞ -algebras, we define the *reduced Hochschild complex* of a functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$ to be the quotient of $C(\Phi)$ given on objects by

$$\overline{C}(\Phi)(m) = \sum_{n \geq 1} \Phi(n+m)/U_n \otimes L_n$$

for $U_n = \sum_{2 \leq i \leq n} \text{Im}(u_i)$, with $u_i : \Phi(n-1+m) \rightarrow \Phi(n+m)$ introducing a unit at the i th position. Lemma 5.4 of [36] shows that the differential of $C(\Phi)$ induces a well-defined differential on $\overline{C}(\Phi)$.

Similarly, for $\Psi : \mathcal{E}^{op} \rightarrow \text{Ch}$ and $m \in \mathbb{N}$, the *reduced coHochschild complex* $\overline{D}(\Psi)$ is the subcomplex of $D(\Psi)$ defined by

$$\overline{D}(\Psi)(m) = \prod_{n \geq 1} \text{Hom}(L_n, K(n) \subset \Psi(n+m))$$

where $K(n) = \bigcap_{i \geq 2}^n \ker(u_i^{op})$. (The kernels $K(n)$ do not define a functor from \mathcal{A}_∞ , so that we cannot define the reduced complex directly as a complex of natural transformations.) Proposition 1.6 below checks that the differential of $D(\Psi)(m)$ restrict to a differential on $\overline{D}(\Psi)(m)$.

Note that $\overline{D}(\Psi)$ is a subfunctor of $D(\Psi) : \mathcal{E}^{op} \rightarrow \text{Ch}$. This follows from the fact that, for $f \in \mathcal{E}^{op}(m, m')$ and any $i \leq n$, we have a commutative diagram

$$\begin{array}{ccc} \Psi(n+m) & \xrightarrow{\Psi(u_i^{op})} & \Psi(n-1+m) \\ \Psi(id_n + f) \downarrow & & \downarrow \Psi(id_{n-1} + f) \\ \Psi(n+m') & \xrightarrow{\Psi(u_i^{op})} & \Psi(n-1+m') \end{array}$$

as u_i^{op} could just as well be written as $u_i^{op} + id_m$ (resp. $u_i^{op} + id_{m'}$). In fact, \overline{D} defines again a self-map of $\text{Fun}(\mathcal{E}, \text{Ch})$.

To prove that $\overline{D}(\Psi)$ is quasi-isomorphic to $D(\Psi)$, we will consider partially reduced Hochschild complexes as well: let

$$\overline{D}_{\leq r}(\Psi)(m) = \prod_{n \geq 1} \text{Hom}(L_n, K(n)_{\leq r} \subset \Psi(n+m))$$

where $K(n)_{\leq r} = \bigcap_{i=2}^{\max(r,n)} \ker(u_i) \subset \Psi(n+m)$. In particular, $\overline{D}_{\leq 1}(\Psi)(m) = D(\Psi)(m)$ and $\overline{D}(\Psi)(m) = \bigcap_{r \geq 1} \overline{D}_{\leq r}(\Psi)(m)$.

Proposition 1.6. *The differential of $D(\Psi)(m)$ restricts to a differential on $\overline{D}(\Psi)(m)$ and on each $\overline{D}_{\leq r}(\Psi)(m)$.*

Proof. We prove the proposition for $\overline{D}(\Psi)(m)$. The same proof applies to each $\overline{D}_{\leq r}(\Psi)(m)$. We write $d_\Psi + d^L$ for the differential of the coHochschild complex, with d^L the coHochschild differential. First note that d_Ψ takes $\overline{D}(\Psi)(m)$ to itself as u_i is of degree 0 so $d_\Psi(\Psi(u_i^{op})(x)) = \Psi(u_i^{op})(d_\Psi x)$. So we are left to consider d^L .

Recall that $d^L = \sum_{n > k} f_{n,k}^{op}$, where each $f_{n,k} = \sum m_{n,k}^i \in \mathcal{A}_\infty(n, k)$. For $x \in K(k) \subset \Psi(k+m)$, we will show that $\Psi(f_{n,k}^{op})(x) = \Psi(f_{n,k}^{op} + id_m)(x) \in K(n) \subset \Psi(n+m)$, i.e. that

each of the sum of terms defined by $f_{n,k}$ take $K(n)$ to $K(n)$. So assume $\Psi(u_i^{op})(x) = 0$ for each $2 \leq i \leq k$. We need to check that $\Psi(u_i^{op})(\Psi(f_{n,k}^{op})(x)) = 0$ for each $2 \leq i \leq n$. For this, we need to consider the composition $f_{n,k} \circ u_i \in \mathcal{A}_\infty^+(n-1, k)$. Recall from [36, Lem 5.4] (which deals with the dual situation) that $f_{n,k} \circ u_i$ has the form $\sum_j \pm(u_{i'} + m_{n-k+1}^j)$ with $2 \leq i' \leq i$ and $i' \leq k$. Hence $(f_{n,k} \circ u_i)^{op} = \sum (f')^{op} \circ u_{i'}^{op}$ with $2 \leq i' \leq i$ maps x to 0 if $x \in K(k)$. \square

Proposition 1.7. *The inclusion $\overline{D}(\Psi)(m) \hookrightarrow D(\Psi)(m)$ is a quasi-isomorphism.*

Proof. (This is an adaptation of the proof of [15, Lem 16]. I would like to apologize in advance about the signs in this proof: they come from the behavior of orientations in the graph complex and they are not very transparent.) The partially reduced complexes $F^r := \overline{D}_{\leq r}(\Psi)(m)$ define a filtration of $D(\Psi)(m)$:

$$\overline{D}(\Psi) = \bigcap_{r \geq 1} F^r \subset \dots \subset F^r \subset F^{r-1} \subset \dots \subset F^1 = D(\Psi).$$

Using the identification $D(\Psi)(m) \cong \prod_{n \geq 1} \Psi(n+m)[1-n]$, we have

$$F^r \cong \prod_{n \geq 1} K(n)_{\leq r}[1-n].$$

Note that for each r , the first $r-1$ factors of F^r and F^{r-1} are identical.

To prove the proposition, we will show that the quotients F^{r-1}/F^r have trivial homology. To identify these quotients, we use the short exact sequence

$$F^r = \prod_{n \geq 1} K(n)_{\leq r}[1-n] \longrightarrow F^{r-1} = \prod_{n \geq 1} K(n)_{\leq r-1}[1-n] \xrightarrow{u} B^r = \prod_{n \geq r} K(n-1)_{\leq r-1}[1-n]$$

with u the 0-map on the factors with $n < r$, and $u = (i^{op})^* \Psi(u_r^{op})$ on the other factors—to ease the notation, we will from now on in the proof simply write u_i^{op} for $(i^{op})^* \Psi(u_r^{op})$ and more generally drop $(i^{op})^* \Psi$ from the notation on morphisms applied to elements of $K(n)$.

Note that u is surjective (the multiplication $m_{r-1,r} \in \mathcal{A}_\infty(n, n-1)$ giving a splitting) and that F^r is indeed the kernel of u . Hence $B^r \cong F^{r-1}/F^r$ as a graded vector space.

We compute the quotient differential on B^r . Recall that for $x = (x_1, x_2, \dots) \in F^{r-1}$, its differential has component n given by $(dx)_n = (-1)^{n-1}(d_\Psi(x_n) - \sum_{k < n} f_{n,k}^{op}(x_k))$, where $f_{n,k} = \sum_{i=1}^n m_{n,k}^i$ where $m_{n,k}^i = \pm m_{n-k+1}^i$ multiplies the entries $i, \dots, i+n-k$. For $y = (y_r, y_{r+1}, \dots) \in B^r$, with each $y_n \in K(n-1)_{\leq r-1}$, define dy by

$$(dy)_n = (-1)^{n-1}(d_\Psi(y_n) + \sum_{k=r}^{n-1} \sum_{i=r}^k (-1)^{n-k}(m_{n-1,k-1}^i)^{op}(y_k))$$

To check that this is the quotient differential, we need to check that the diagram

$$\begin{array}{ccc} F^{r-1} & \xrightarrow{u} & B^r \\ \downarrow d & & \downarrow d \\ F^{r-1} & \xrightarrow{u} & B^r \end{array}$$

commutes. Let $x = (x_1, x_2, \dots) \in F^{r-1}$. Then for $n \geq r$,

$$\begin{aligned} (u \circ d(x))_n &= (-1)^{n-1}(u_r^{op}(d_\Psi(x_n)) - \sum_{k < n} u_r^{op}(f_{n,k}^{op}(x_k))) \\ &= (-1)^{n-1}(u_r^{op}(d_\Psi(x_n)) - \sum_{k < n} (f_{n,k} \circ u_r)^{op}(x_k)). \end{aligned}$$

We have $f_{n,k} \circ u_r = \sum_{i=1}^n m_{n,k}^i \circ u_r$. There are 3 possibilities: r can be before the multiplication, at the multiplication, or after it. If $n - k > 1$, and r is at the multiplication, i.e. $i \leq r \leq i + n - k$ (understood cyclically), then the composition is 0. If $n - k = 1$, there are exactly two multiplications which include r , namely $i = r - 1$ and $i = r$ (again understood cyclically, i.e. mod n), and the two terms will cancel each other. Hence there will be no contribution of that form. If r is after the multiplication, $m_{n,k}^i \circ u_r = u_j \circ m_{n-1,k-1}^i$ for some $j < r$, and hence it acts as 0 on F^{r-1} . More precisely, this happens when $r > i + n - k$ cyclically, except when $i + n - k = 1$. Hence the only the terms contributing to this sum are those with $i > r$ and $i + n - k < n + 2$ (now non-cyclically), using the fact that $r > 1$ for the case $i + n - k = n + 1$. This means that only the terms $m_{n,k}^i$ with $r < i < k + 2$ contribute, which in particular requires $k \geq r$. Now in these cases we have $m_{n,k}^i \circ u_r = (-1)^{n-k} u_r \circ m_{n-1,k-1}^{i-1}$. So

$$(u \circ d(x))_n = (-1)^{n-1} (d_\Psi(u_r^{op}(x_n))) + \sum_{k=r}^{n-1} \sum_{i=1}^k (-1)^{n-k} (m_{n-1,k-1}^{i-1})^{op}(u_r^{op}(x_k))$$

which is exactly the n th component of $d \circ u(x)$.

We now show that B^r has trivial homology. Define $s : B^r \rightarrow B^r$ of degree $+1$ by $s(y)_n = (-1)^{n+r} u_r^{op}(y_{n+1})$. Explicitly, we have

$$\begin{aligned} (sd(y))_n &= (-1)^{n+r} u_r^{op}(dy)_{n+1} \\ &= (-1)^r (u_r^{op}(d_\Psi(y_{n+1}))) + \sum_{k=r}^n \sum_{i=r}^k (-1)^{n-k+1} u_r^{op}((m_{n,k-1}^i)^{op}(y_k)) \\ (ds(y))_n &= (-1)^{n-1} ((-1)^{n+r} d_\Psi(u_r^{op}(y_{n+1}))) \\ &\quad + \sum_{k'=r}^{n-1} \sum_{i'=r}^{k'} (-1)^{n-k'+r+k'} (m_{n-1,k'-1}^{i'})^{op}(u_r^{op}(y_{k'+1})) \end{aligned}$$

First note that for $k = n$ and $i = r$ in the first line, we have $u_r^{op}((m_{n,n-1}^r)^{op}(y_n)) = (m_{n,n-1}^r \circ u_r)^{op} y_n = (-1)^{r+1} y_n$. We would like to show that no other term contribute to the sum $(sd(y))_n + (ds(y))_n$.

If $i = r$ in the first sum with $k < n$, the composition $m_{n,k-1}^r \circ u_r = 0$, so such terms do not contribute. In particular, as $k = r$ implies $i = r$, there are no contribution of that form.

In the first sum, we are thus left with the terms $r < k \leq n$ and $r < i \leq k$. Then the composition $m_{n,k-1}^i \circ u_r = (-1)^{n-k+1} u_r \circ m_{n-1,k-2}^{i-1}$. Choosing $k' = k - 1$ and $i' = i - 1$ gives an identification between those terms and minus all the terms of the second sum.

Hence $sd + ds = id$. It follows that B^r has trivial homology and thus that each inclusion $F^r \hookrightarrow F^{r-1}$ is a quasi-isomorphism. To obtain the desired quasi-isomorphism, we consider the short exact sequence

$$\overline{D}(\Psi) = \cap_r F^r \rightarrow D(\Psi) = F^1 \rightarrow \cup_r B^r.$$

As $H_*(B^r) = 0$ for each r , we have that $H_*(\cup_r B^r) = 0$ (any cycle being in some B^r), which implies the result. \square

Proposition 1.8. *The quotient map $C(\Phi) \rightarrow \overline{C}(\Phi)$ is a quasi-isomorphism of functors.*

Proof. The proof is a dual version of the previous proof: we consider the partial quotients of $C(\Phi)(m) \cong \oplus_{n \geq 1} \Phi(n + m)$

$$F_r = \oplus_{n \geq 1} \Phi(n + m) / U_{\leq r} \quad \text{where} \quad U_{\leq r} = \sum_{i=2}^{\max(n,r)} Im(u_i)$$

Then we have a sequence of quotient maps

$$C(\Phi)(m) = F_1 \rightarrow F_2 \rightarrow \cdots \rightarrow F_\infty = \overline{C}(\Phi)(m).$$

We compute the quotients F_r/F_{r-1} and $F_\infty/F_1 = \overline{C}(\Phi)(m)/C(\Phi)(m)$ via short exact sequences

$$A_r \xrightarrow{u_r} F_{r-1} \rightarrow F_r \quad \text{and} \quad A_\infty = \cup_r A_r \rightarrow F_1 \rightarrow F_\infty.$$

Again, we want to show that the A_r 's are acyclic. One checks that, as a graded vector space, $A_r = \bigoplus_{n \geq r} \Phi(n-1+m)[n-1]$ and that the quotient differential is defined on $x \in \Phi(n-1+m)$ by

$$dx = d_\Phi(x) + (-1)^{|x|} \sum_{k=r}^{n-1} \sum_{i=r}^k (-1)^{n-k} m_{n-1, k-1}^i(x).$$

Then define $s : A_r \rightarrow A_r$ by $s(x) = (-1)^{|x|} u_r(x)$ if $x \in \Phi(n-1+m)$. This satisfies $sd + ds = id$, which shows that each A_r is acyclic, and hence that their union is acyclic. \square

2. FORMAL AND NATURAL OPERATIONS ON THE HOCHSCHILD COMPLEX

In this section, we first give a model for the formal operations $\text{Nat}_\mathcal{E}$, the natural operations on the Hochschild complex of *generalized \mathcal{E} -algebras*, i.e. functors $\mathcal{E} \rightarrow \text{Ch}$ without any monoidality assumptions. In Section 2.2 we then compare this chain complex to the chain complex of natural operations $\text{Nat}_\mathcal{E}^\otimes$ on the Hochschild complex of actual algebras. Algebras here are algebras over an arbitrary prop with \mathcal{A}_∞ -multiplication (\mathcal{E}, i) . We say that \mathcal{E} has units if $i : \mathcal{A}_\infty \rightarrow \text{Ch}$ extends to a functor $i^+ : \mathcal{A}_\infty^+ \rightarrow \text{Ch}$.

2.1. Formal operations. For each $m, n \geq 0$, we have the dg-functor

$$C_\mathcal{E}^{(n,m)} : \text{Fun}(\mathcal{E}, \text{Ch}) \longrightarrow \text{Ch}$$

taking a functor Φ to the chain complex $C^n(\Phi)(m)$ of Definition 1.2, the iterated Hochschild complex of Φ evaluated at m . Consider the category $\text{Nat}_\mathcal{E}$ with objects pairs of natural numbers $\begin{bmatrix} n \\ m \end{bmatrix}$ with $n, m \geq 0$, and morphisms the Hom-complexes of the above functors:

$$\text{Nat}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}) := \text{Hom}(C_\mathcal{E}^{(n_1, m_1)}, C_\mathcal{E}^{(n_2, m_2)}).$$

Similarly, when \mathcal{E} has units, we define $\overline{\text{Nat}}_\mathcal{E}$ as the category with morphisms the Hom-complexes of the iterated reduced Hochschild complexes of Section 1.2:

$$\overline{\text{Nat}}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}) := \text{Hom}(\overline{C}_\mathcal{E}^{(n_1, m_1)}, \overline{C}_\mathcal{E}^{(n_2, m_2)}).$$

The representable functors $\mathcal{E}(p, -) : \mathcal{E} \rightarrow \text{Ch}$ have a well-defined Hochschild, and n_1 -iterated Hochschild complex $C^{n_1}(\mathcal{E}(p, -))$. By [36, Prop 5.5], these functors for varying p 's assemble to a define functor

$$C^{n_1} \mathcal{E}(-, -) : \mathcal{E}^{op} \times \mathcal{E} \longrightarrow \text{Ch}$$

so that it makes sense to subsequently take an n_2 -iterated coHochschild complex construction in the first variable to obtain a new functor

$$D^{n_2} C^{n_1} \mathcal{E}(-, -) : \mathcal{E}^{op} \times \mathcal{E} \longrightarrow \text{Ch}.$$

Theorem 2.1. *Let (\mathcal{E}, i) be a prop with \mathcal{A}_∞ -multiplication. Then there are isomorphisms of chain complexes*

$$\text{Nat}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}) \cong D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2)$$

and, when \mathcal{E} has units,

$$\overline{\text{Nat}}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}) \cong \overline{D}^{n_1} \overline{C}^{n_2} \mathcal{E}(-, -)(m_1, m_2).$$

Moreover in that case we have $\overline{\text{Nat}}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}) \simeq \text{Nat}_\mathcal{E}(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix})$.

More explicitly, this says that

$$\mathrm{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) \cong \prod_{j_1, \dots, j_{n_1} \geq 1} \bigoplus_{k_1, \dots, k_{n_2} \geq 1} \mathcal{E}(j + m_1, k + m_2) [k - j + n_1 - n_2]$$

with $j = j_1 + \dots + j_{n_1}$, $k = k_1 + \dots + k_{n_2}$, and where the square brackets indicate the degree shift. The differential is the sum of the differential of \mathcal{E} and the Hochschild and coHochschild differentials. Explicitly, in the case $n_1 = 1 = n_2$ for simplicity, an element $\underline{g} \in \mathrm{Nat}_{\mathcal{E}}([m_1]^{1}, [m_2]^{1})$ under the above isomorphism has the form $\underline{g} = \{g_j\}_{j \geq 1}$ with $g_j = g_{j, k_1} + \dots + g_{j, k_{r_j}}$ and $g_{j, k_i} \in \mathcal{E}(j + m_1, k_i + m_2)$. Its differential has \underline{j} th component $d(\underline{g})_{\underline{j}}$ given by

$$(-1)^{j-1} \left(d_{\mathcal{E}}(g_j) + \sum_{i=1}^{r_j} \sum_{k=1}^{k_i-1} (-1)^{|g_{j, k_i}|} (f_{k_i, k} + id_{m_2}) \circ g_{j, k_i} - \sum_{j'=1}^{j-1} g_{j'} \circ (f_{j, j'} + id_{m_1}) \right).$$

We say that a functor $\mathcal{E} \rightarrow \mathcal{E}'$ is a *quasi-isomorphism of categories* if it is the identity on objects and it induces a quasi-isomorphism on each morphism complex.

Using the homotopy invariance of the Hochschild and coHochschild constructions (Proposition 1.5 and [36, Prop 5.6]), we immediately get the following corollary:

Corollary 2.2. *A quasi-isomorphism of categories $\mathcal{E} \xrightarrow{\simeq} \mathcal{E}'$ induces a quasi-isomorphism $\mathrm{Nat}_{\mathcal{E}} \xrightarrow{\simeq} \mathrm{Nat}_{\mathcal{E}'}$.*

To prove Theorem 2.1, we first show that the above chain complex does act by natural transformations on the Hochschild complex of generalized \mathcal{E} -algebras.

Lemma 2.3. *For any $\Phi : \mathcal{E} \rightarrow \mathrm{Ch}$, there is a chain map*

$$C^{n_1}(\Phi)(m_1) \otimes D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2) \longrightarrow C^{n_2}(\Phi)(m_2)$$

which is natural in Φ . This map restricts to a map

$$\overline{C}^{n_1}(\Phi)(m_1) \otimes \overline{D}^{n_1} \overline{C}^{n_2} \mathcal{E}(-, -)(m_1, m_2) \longrightarrow \overline{C}^{n_2}(\Phi)(m_2)$$

in the reduced case.

Proof. To simplify notations write $\underline{j} = (j_1, \dots, j_{n_1})$, $j = j_1 + \dots + j_{n_1}$ and $L_{\underline{j}} = L_{j_1} \otimes \dots \otimes L_{j_{n_1}}$, and similarly for $\underline{k} = (k_1, \dots, k_{n_2})$. In the unreduced case, the action is defined as follows:

$$\begin{aligned} & (\Phi(- + m_1) \otimes_{\mathcal{A}_{\infty}} \mathcal{L}) \otimes \mathrm{Hom}_{\mathcal{A}_{\infty}^{op}}(\mathcal{L}, \mathcal{E}(- + m_1, - + m_2) \otimes_{\mathcal{A}_{\infty}} \mathcal{L}) \\ & \quad \downarrow \\ & \Phi(- + m_1) \otimes_{\mathcal{A}_{\infty}} \mathcal{E}(- + m_1, - + m_2) \otimes_{\mathcal{A}_{\infty}} \mathcal{L} \\ & \quad \downarrow \\ & \Phi(- + m_2) \otimes_{\mathcal{A}_{\infty}} \mathcal{L} \end{aligned}$$

where the first arrow applies the homomorphism to the $\mathcal{A}_{\infty}^{op}$ -module \mathcal{L} used to define the Hochschild complex in Section 1, and the second arrow is induced by the action of \mathcal{E} on Φ . As the action on Φ is via its \mathcal{E} -structure, it is natural with respect to natural transformations of functors $\Phi \rightarrow \Phi'$.

More explicitly, the action is the composition

$$\begin{aligned} & (\bigoplus_{\underline{j}} \Phi(j + m_1) \otimes L_{\underline{j}}) \otimes \prod_{\underline{j}} \mathrm{Hom}(L_{\underline{j}}, \bigoplus_{\underline{k}} \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}}) \\ & \quad \downarrow \\ & \bigoplus_{\underline{j}, \underline{k}} \Phi(j + m_1) \otimes \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}} \\ & \quad \downarrow \\ & \bigoplus_{\underline{k}} \Phi(k + m_2) \otimes L_{\underline{k}} \end{aligned}$$

which restricts to a well-defined action of $\overline{D}^{n_1} \overline{C}^{n_2} \mathcal{E}(-, -)(m_1, m_2)$ in the reduced case because the reduced coHochschild complex only involves homomorphisms with image in the kernel of precomposition by the maps u_i whose images are quotiented out in the reduced Hochschild complex of Φ . \square

This lemma gives a map

$$F : D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2) \rightarrow \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2})$$

(and reduced version) which is easily seen to be injective using as Φ 's the representable functors $\mathcal{E}(p, -)$. The composition on the complexes $D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2)$ for varying m_i 's and n_i 's which corresponds to the composition of natural transformations in $\text{Nat}_{\mathcal{E}}$ under this map is of the form

$$\begin{aligned} & \prod_j \text{Hom}(L_j, \bigoplus_k \mathcal{E}(j, k) \otimes L_k) \otimes \prod_k \text{Hom}(L_k, \bigoplus_l \mathcal{E}(k, l) \otimes L_l) \\ & \quad \downarrow \\ & \prod_j \text{Hom}(L_j, \bigoplus_{k,l} \mathcal{E}(j, k) \otimes \mathcal{E}(k, l) \otimes L_l) \\ & \quad \downarrow \\ & \prod_j \text{Hom}(L_j, \bigoplus_l \mathcal{E}(j, l) \otimes L_l). \end{aligned}$$

Proof of Theorem 2.1. We will construct a pointwise inverse to the map F given by the lemma above. For each $m_1, m_2, n_1, n_2 \geq 0$, a map

$$\begin{aligned} G : \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) & \longrightarrow D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2) \\ & = \prod_{\underline{j}} \text{Hom}(L_{\underline{j}}, \bigoplus_{\underline{k}} \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}}) \end{aligned}$$

is determined by giving, for each natural transformation ν and each tuple $\underline{j} = (j_1, \dots, j_{n_1})$, an element $G(\nu)_{\underline{j}}(l_{\underline{j}}) \in \bigoplus_{\underline{k}} \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}}$, for $l_{\underline{j}}$ the generator of $L_{\underline{j}}$. To produce such an element given tuple $\underline{j} = (j_1, \dots, j_{n_1})$, we consider the action of $\text{Nat}_{\mathcal{E}}$ on the Hochschild complex of the functor $\mathcal{E}(j + m_1, -)$ for $j = j_1 + \dots + j_{n_1}$:

$$\begin{array}{ccc} C^{n_1}(\mathcal{E}(j + m_1, -))(m_1) \otimes \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) & \xrightarrow{\alpha} & C^{n_2}(\mathcal{E}(j + m_1, -))(m_2) \\ \downarrow = & & \uparrow = \\ (\bigoplus_{\underline{j}'} \mathcal{E}(j + m_1, j' + m_1) \otimes L_{\underline{j}'}) \otimes \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) & \longrightarrow & \bigoplus_{\underline{k}} \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}}. \end{array}$$

We define

$$G(\nu)_{\underline{j}}(l_{\underline{j}}) := \nu(id_{j+m_1} \otimes l_{\underline{j}}) = \alpha((id_{j+m_1} \otimes l_{\underline{j}}) \otimes \nu).$$

We check that G is a chain map: as α is a chain map, we have that $G(d\nu)_{\underline{j}}(l_{\underline{j}}) = (-1)^{j-1}(d(\nu(id_{j+m_1} \otimes l_{\underline{j}})) - \nu(d(id_{j+m_1} \otimes l_{\underline{j}})))$. The first term gives exactly the differential of the Hochschild complex of \mathcal{E} , so we are left to identify the second term with the coHochschild differential. We have $d(id_{j+m_1} \otimes l_{\underline{j}}) = \sum_{\underline{j}' < \underline{j}} f_{\underline{j}, \underline{j}'} \otimes l_{\underline{j}'}$, where $\underline{j}' < \underline{j}$ are the tuples (j'_1, \dots, j'_{n_1}) with $j'_i = j_i$ except for one $i = s$ where $j'_s < j_s$, and $f_{\underline{j}, \underline{j}'} = (-1)^{j_1 + \dots + j_{s-1} - s + 1} id + \dots + f_{j_s, j'_s} + \dots + id$. By naturality, $\nu(\sum_{\underline{l}} f_{\underline{j}, \underline{l}} \otimes l_{\underline{l}}) = \sum_{\underline{l}} (\nu(id_{l+m_1} \otimes l_{\underline{l}}) \circ (f_{\underline{j}, \underline{l}} + id_{m_1}))$, which is the coHochschild differential.

One easily checks that $G \circ F$ is the identity on $D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2)$: an element in $g \in \prod_{\underline{j}} \text{Hom}(L_{\underline{j}}, \bigoplus_{\underline{k}} \mathcal{E}(j + m_1, k + m_2) \otimes L_{\underline{k}})$ defines a natural transformation via the map F which takes $id_{j+m_1, -} \otimes l_{\underline{j}}$ to its own j th component $g_{\underline{j}}$. To check that $F \circ G$ is the identity on $\text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2})$, we need to check that the natural transformation $F \circ G(\nu)$ acts the same way as ν on any functor $\Phi : \mathcal{E} \rightarrow \text{Ch}$. Consider an element $x \otimes l_{\underline{j}} \in C^{n_1}(\Phi)(m_1) = \bigoplus_{\underline{j}} \Phi(j + m_1) \otimes L_{\underline{j}}$. There is an associated natural transformation $\nu_x : \mathcal{E}(j + m_1, -) \rightarrow \Phi$ defined by evaluation at x which takes id_{j+m_1} to x . By naturality

of the natural transformations, we have a commutative diagram

$$\begin{array}{ccc} C^{n_1}(\mathcal{E}(j+m_1, -))(m_1) \otimes \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) & \xrightarrow{\alpha} & C^{n_2}(\mathcal{E}(j+m_1, -))(m_2) \\ \downarrow & & \downarrow \\ C^{n_1}(\Phi)(m_1) \otimes \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) & \xrightarrow{\alpha} & C^{n_2}(\Phi)(m_2) \end{array}$$

where the vertical arrows are induced by the natural transformations. By definition, the actions of ν and $F \circ G(\nu)$ are the same on the elements of the type $id_{j+m_1} \otimes l_j$, and thus they agree on $x \otimes l_j$ by the commutativity of the above diagram.

Note again that the inverse G restrict to an inverse in the reduced case.

For the quasi-isomorphism between the reduced and unreduced cases, we use the sequence of quasi-isomorphisms

$$\begin{array}{ccc} \overline{D}^{n_1} \overline{C}^{n_2} \mathcal{E}(-, -)(m_1, m_2) & \xrightarrow{\simeq} & D^{n_1} \overline{C}^{n_2} \mathcal{E}(-, -)(m_1, m_2) \\ & & \uparrow \simeq \\ & & D^{n_1} C^{n_2} \mathcal{E}(-, -)(m_1, m_2) \end{array}$$

given by first applying Proposition 1.7 iteratively, together with Proposition 1.5, and then Proposition 1.8 together with [36, Prop 4.5]. \square

Remark 2.4. The category $\text{Nat}_{\mathcal{E}}$ is an *extension of the Hochschild core category of \mathcal{E}* in the sense of [36, Sec 5.2] as it satisfies that

$$\text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) \cong C^{n_2}(\mathcal{E}(m_1, -))(m_2).$$

Now by Corollary 5.12 of [36], any extension $\tilde{\mathcal{E}}$ of \mathcal{E} admits a functor $\tilde{\mathcal{E}} \rightarrow \text{Nat}_{\mathcal{E}}$ as it acts by natural transformations on the Hochschild complex of functors $\Phi : \mathcal{E} \rightarrow \text{Ch}$. Another characterization of $\text{Nat}_{\mathcal{E}}$ is that it is the universal extension of \mathcal{E} .

2.2. Restriction to natural operations. Recall from the introduction the category $\text{Fun}^{\otimes}(\mathcal{E}, \text{Ch})$ of \mathcal{E} -algebras whose objects are strong symmetric monoidal dg-functors $\Phi : \mathcal{E} \rightarrow \text{Ch}$, i.e. with natural isomorphisms $\Phi(n) \otimes \Phi(m) \xrightarrow{\cong} \Phi(n+m)$ compatible with the symmetries in \mathcal{E} , and whose morphisms are natural transformation $\theta : \Phi \rightarrow \Psi$ satisfying

$$\theta_n = (\theta_1)^{\otimes n} : \Phi(n) \cong \Phi(1)^{\otimes n} \longrightarrow \Psi(n) \cong \Psi(1)^{\otimes n}$$

with $\theta_1 : \Phi(1) \rightarrow \Psi(1)$ a chain map.

Writing $C_{\mathcal{E}}^{\otimes(n,m)}$ for the functor taking Φ to $(C(\Phi)(0))^{\otimes n} \otimes (\Phi(1))^{\otimes m}$, Proposition 5.10 of [36] says that the diagram

$$\begin{array}{ccc} \text{Fun}(\mathcal{E}, \text{Ch}) & \xrightarrow{C_{\mathcal{E}}^{(n,m)}} & \text{Ch} \\ \uparrow & \searrow C_{\mathcal{E}}^{\otimes(n,m)} & \\ \text{Fun}^{\otimes}(\mathcal{E}, \text{Ch}) & & \end{array}$$

commutes. Note that $\text{Fun}^{\otimes}(\mathcal{E}, \text{Ch})$ is not a dg-category as such, but it can be replaced, if one wants to stay in the world of dg-categories, by the sub-dg-category of $\text{Fun}(\mathcal{E}, \text{Ch})$ it generates, that is allowing sums of monoidal natural transformations even when these are not themselves monoidal. Let

$$\text{Nat}_{\mathcal{E}}^{\otimes}([m_1]^{n_1}, [m_2]^{n_2}) := \text{Hom}(C_{\mathcal{E}}^{\otimes(n_1, m_1)}, C_{\mathcal{E}}^{\otimes(n_2, m_2)})$$

denote the chain complex of natural transformations of these functors. There is a restriction map

$$r : \text{Nat}_{\mathcal{E}}([m_1]^{n_1}, [m_2]^{n_2}) \longrightarrow \text{Nat}_{\mathcal{E}}^{\otimes}([m_1]^{n_1}, [m_2]^{n_2})$$

as a natural transformation on all functors gives in particular a natural transformation on the subcategory of symmetric monoidal functors. We will show in this section that the restriction map r is an isomorphism provided that \mathcal{E} is “complete” in a sense defined below.

Recall that $\text{Nat}_{\mathcal{E}}([m_1], [m_2]) \cong \mathcal{E}(m_1, m_2)$. On the other hand, $\text{Nat}_{\mathcal{E}}^{\otimes}([m_1], [m_2]) = \text{Hom}(U^{\otimes m_1}, U^{\otimes m_2})$ for $U : \text{Fun}^{\otimes}(\mathcal{E}, \text{Ch}) \rightarrow \text{Ch}$ the forgetful functor from the category of \mathcal{E} -algebras to the category of chain complexes taking Φ to $\Phi(1)$, and $U^{\otimes m_i}$ its tensor powers. So when $n_1 = 0 = n_2$, the restriction map r above is the representation map of the prop \mathcal{E} , i.e. the map

$$\rho : \mathcal{E}(m_1, m_2) \longrightarrow \text{Hom}(U^{\otimes m_1}, U^{\otimes m_2})$$

which associates to a morphism of \mathcal{E} the operation it induces on all \mathcal{E} -algebras.

Definition 2.5. *We say that a prop \mathcal{E} is complete if the representation functor ρ defined above is an isomorphism. If \mathcal{E} is not complete, we call the target of ρ , the prop of natural transformations of the forgetful functors, its completion, and denote it $\widehat{\mathcal{E}}$.*

So given a prop \mathcal{E} , its completion $\widehat{\mathcal{E}}$ is a prop defined by

$$\widehat{\mathcal{E}}(m_1, m_2) = \text{Hom}(U^{\otimes m_1}, U^{\otimes m_2}) = \text{Nat}_{\mathcal{E}}^{\otimes}([m_1], [m_2])$$

with U as above.

Lemma 2.6. *Let \mathcal{E} be a prop and $\widehat{\mathcal{E}}$ its completion. Then their categories of algebras are isomorphic: $\text{Fun}^{\otimes}(\mathcal{E}, \text{Ch}) \cong \text{Fun}^{\otimes}(\widehat{\mathcal{E}}, \text{Ch})$.*

Proof. The representation map ρ above induces a functor

$$\rho^* : \text{Fun}^{\otimes}(\widehat{\mathcal{E}}, \text{Ch}) \longrightarrow \text{Fun}^{\otimes}(\mathcal{E}, \text{Ch})$$

by precomposition. On the other hand, any \mathcal{E} -algebra is an $\widehat{\mathcal{E}}$ -algebra by definition of $\widehat{\mathcal{E}}$, hence this functor is the identity on objects. Now morphisms of $\mathcal{E}/\widehat{\mathcal{E}}$ -algebras are chain maps $A \rightarrow B$ that commute with the maps in $\mathcal{E}/\widehat{\mathcal{E}}$. This implies directly that the functor is injective on morphisms as both $\mathcal{E}(A, B)$ and $\widehat{\mathcal{E}}(A, B)$ are subsets of $\text{Hom}(A, B)$ and the functor commutes with the inclusions. Surjectivity of ρ^* follows from the definition of $\widehat{\mathcal{E}}$: the morphisms in $\widehat{\mathcal{E}}$ are precisely the natural transformations that commute with all maps of \mathcal{E} -algebras, so maps of \mathcal{E} -algebras are also maps of $\widehat{\mathcal{E}}$ -algebras. \square

As the completion of a prop is defined in terms of its category of algebras $\text{Fun}^{\otimes}(\mathcal{E}, \text{Ch})$, the lemma immediately implies the following

Corollary 2.7. *The completion of a prop is complete.*

Similarly, when \mathcal{E} is a prop with \mathcal{A}_{∞} -multiplication, we get

Corollary 2.8. *Let (\mathcal{E}, i) be a prop with \mathcal{A}_{∞} -multiplication and $(\widehat{\mathcal{E}}, \rho \circ i)$ its completion. Then*

$$\text{Nat}_{\mathcal{E}}^{\otimes}([m_1], [m_2]) \cong \text{Nat}_{\widehat{\mathcal{E}}}^{\otimes}([m_1], [m_2])$$

We are now ready to prove the main result of this section, which says that the injectivity and surjectivity of the restriction map r is determined by these conditions on the representation map ρ :

Theorem 2.9. *Let (\mathcal{E}, i) be a prop with \mathcal{A}_{∞} -multiplication. The restriction map $r : \text{Nat}_{\mathcal{E}} \longrightarrow \text{Nat}_{\widehat{\mathcal{E}}}^{\otimes}$ is an isomorphism if and only if \mathcal{E} is complete.*

More precisely, let $(\widehat{\mathcal{E}}, \rho \circ i)$ denote the completion of \mathcal{E} . Then r is injective (resp. surjective) if and only if the map $\rho : \mathcal{E} \rightarrow \widehat{\mathcal{E}}$ is injective (resp. surjective).

In particular, we can always write the natural operations on the Hochschild complex of \mathcal{E} -algebras in terms of the generalized natural operations for its completion $\widehat{\mathcal{E}}$:

Corollary 2.10. *Let (\mathcal{E}, i) be a prop with \mathcal{A}_∞ -multiplication and $(\widehat{\mathcal{E}}, \rho \circ i)$ its completion. Then*

$$\mathrm{Nat}_{\mathcal{E}}^{\otimes}([m_1], [m_2]) \cong \mathrm{Nat}_{\widehat{\mathcal{E}}}([m_1], [m_2])$$

This follows from Theorem 2.9 and from Corollary 2.8.

Proof of Theorem 2.9. Let $\nu \in \mathrm{Nat}_{\mathcal{E}}([m_1], [m_2])$ and suppose that $r(\nu) = 0$. Then for any \mathcal{E} -algebra $\Phi \in \mathrm{Fun}^{\otimes}(\mathcal{E}, \mathrm{Ch})$, the map $\nu_{\Phi} : C^{m_1}(\Phi)(m_1) \rightarrow C^{m_2}(\Phi)(m_2)$ is the 0-map. We have

$$\nu = (\nu_{\underline{j}}) \in \prod_{\underline{j}} \bigoplus_{\underline{k}} \mathrm{Hom}(L_{\underline{j}}, \mathcal{E}(j+m_1, k+m_2) \otimes L_{\underline{k}}) \cong \prod_{\underline{j}} \bigoplus_{\underline{k}} \mathcal{E}(j+m_1, k+m_2)[k-j+n_1-n_2]$$

and ν_{Φ} is defined on $\Phi(j+m_1) \otimes L_{\underline{j}} \subset C^{m_1}(\Phi)(m_1)$ as the composition

$$\begin{array}{ccc} \Phi(j+m_1) \otimes L_{\underline{j}} & \xrightarrow{\nu_{\underline{j}}} & \bigoplus_{\underline{k}} \Phi(j+m_1) \otimes \mathcal{E}(j+m_1, k+m_2) \otimes L_{\underline{k}} \\ & \searrow \nu_{\Phi} & \downarrow \\ & & \bigoplus_{\underline{k}} \Phi(k+m_2) \otimes L_{\underline{k}} \end{array}$$

By the above diagram, ν_{Φ} acting as 0 for each Φ is equivalent to each $\nu_{\underline{j}}(l_{\underline{j}})_{\underline{k}} \in \mathcal{E}(j+m_1, k+m_2)$ acting as 0 for each Φ , i.e. $\rho(\nu_{\underline{j}}) = 0$ for each \underline{j} . But that implies that each $\nu_{\underline{j}}(l_{\underline{j}})_{\underline{k}} = 0$ when ρ is injective, and hence that $\nu = 0$, which proves the injectivity part of the statement.

Consider now $\nu \in \mathrm{Nat}_{\mathcal{E}}^{\otimes}([m_1], [m_2])$. Then for each \mathcal{E} -algebra Φ , we have a map

$$\nu_{\Phi} : \bigoplus_{\underline{j}} \Phi(j+m_1) \otimes L_{\underline{j}} \longrightarrow \bigoplus_{\underline{k}} \Phi(k+m_2) \otimes L_{\underline{k}}.$$

This map is given by a collection of maps $(\nu_{\Phi})_{(\underline{j}, \underline{k})} : \Phi(j+m_1) \rightarrow \Phi(k+m_2)$ which are natural in Φ . In other words these maps define an element $(\nu)_{(\underline{j}, \underline{k})} \in \widehat{\mathcal{E}}(j+m_1, k+m_2)$. If $\rho : \mathcal{E} \rightarrow \widehat{\mathcal{E}}$ is surjective, this implies that $(\nu)_{(\underline{j}, \underline{k})} = \rho(f)$ for some $f \in \mathcal{E}(j+m_1, k+m_2)$, and we have that $(\nu_{\Phi})_{(\underline{j}, \underline{k})} = \Phi(f)$ for each Φ . Hence ν comes from an element in $\mathrm{Nat}_{\mathcal{E}}([m_1], [m_2])$. \square

The representation map $\rho : \mathcal{E} \rightarrow \widehat{\mathcal{E}}$ is injective exactly when any two morphisms in \mathcal{E} can be distinguished by their actions on \mathcal{E} -algebras, i.e. if for every j, k and $f \neq g \in \mathcal{E}(j, k)$, there exists a $\Phi \in \mathrm{Fun}^{\otimes}(\mathcal{E}, \mathrm{Ch})$ such that $\Phi(f) \neq \Phi(g)$. This is always possible when \mathcal{E} admits free algebras, which is the case when \mathcal{E} is the prop associated to an operad:

Example 2.11. Suppose $\mathcal{E}_{\mathcal{P}}$ is the prop associated to an operad $\mathcal{P} = \{\mathcal{P}(n)\}_{n \geq 0}$, so

$$\mathcal{E}_{\mathcal{P}}(p, q) = \bigoplus_{n_1 + \dots + n_q = p} \mathcal{P}(n_1) \otimes \dots \otimes \mathcal{P}(n_q) \otimes_{\Sigma_{n_1} \times \dots \times \Sigma_{n_q}} \mathbb{Z}\Sigma_p.$$

Let $\widehat{\mathcal{E}}_{\mathcal{P}}$ denote its completion. Then the representation map $\rho : \mathcal{E}_{\mathcal{P}} \rightarrow \widehat{\mathcal{E}}_{\mathcal{P}}$ is injective as any two elements of $\mathcal{E}_{\mathcal{P}}$ can be distinguished by their actions on some free algebra $\mathcal{P}(X) = \bigoplus_n \mathcal{P}(n) \otimes_{\Sigma_n} X^{\otimes n}$.

The representation map is however not injective in general, as shown by the following example:

Example 2.12. (lunch time example of Oscar Randal-Williams.) Let Frob be the prop of commutative Frobenius algebras (i.e. the prop of (closed) 2-dimensional topological quantum field theories, see e.g. [20, 1.2]) over a field. Consider the quotient $\mathrm{Frob}/_{(T=0)}$ of Frob by the relation that the operation associated to the torus equals the 0-map.

As the torus operation in the Frobenius prop is multiplication by the dimension of the algebra (see [20, 1.2.29]), this means that only a 0-dimensional Frobenius algebra can be an algebra over this prop. Hence there is only one $\text{Frob}/_{(T=0)}$ -algebra, the trivial 0-dimensional one, and elements in the prop cannot be distinguished by their action on algebras.

Surjectivity of the representation map is more difficult to check. It is at least satisfied in characteristic 0 by props associated to operads concentrated in degree 0:

Example 2.13 (Fresse). Let k be a field of characteristic 0 and \mathcal{P} be an operad in Vect_k , the category of vector spaces over k . Let $\mathcal{E}_{\mathcal{P}}$ denote its associated prop as in Example 2.11. Then $\mathcal{E}_{\mathcal{P}}$ is complete in Vect_k , i.e. $\mathcal{E}_{\mathcal{P}}(r, s) \cong \widehat{\mathcal{E}}_{\mathcal{P}}(r, s) := \text{Hom}((U_V)^{\otimes r}, (U_V)^{\otimes s})$ for $U_V : \text{Fun}^{\otimes}(\mathcal{E}_{\mathcal{P}}, \text{Vect}_k) \rightarrow \text{Vect}_k$ the forgetful functor from \mathcal{P} -algebras to vector spaces. Indeed, consider the functors

$$\begin{aligned} id^{\otimes r} : \text{Vect}_k &\longrightarrow \text{Vect}_k &: X &\mapsto X^{\otimes r} \\ \mathcal{P}(-)^{\otimes s} : \text{Vect}_k &\longrightarrow \text{Vect}_k &: X &\mapsto \mathcal{P}(X)^{\otimes s} = \left(\bigoplus_n \mathcal{P}(n) \otimes_{\Sigma_n} X^{\otimes n} \right)^{\otimes s}. \end{aligned}$$

Let $\text{Hom}(id^{\otimes r}, \mathcal{P}(-)^{\otimes s})$ denote the natural transformations between these functors. There is map

$$\mathcal{E}_{\mathcal{P}}(r, s) = \bigoplus_{r_1 + \dots + r_s = r} \mathcal{P}(r_1) \otimes \dots \otimes \mathcal{P}(r_s) \otimes_{\Sigma_{r_1} \times \dots \times \Sigma_{r_s}} \mathbb{Z}\Sigma_r \xrightarrow{\eta} \text{Hom}(id^{\otimes r}, \mathcal{P}(-)^{\otimes s})$$

mapping a morphism $a = a_1 \otimes \dots \otimes a_s \otimes \sigma$ to the natural transformation ν_a defined on a vector space X as the composition

$$\nu_a(X) : X^{\otimes r} \xrightarrow{\sigma} X^{\otimes r} \hookrightarrow (a_1 \otimes_{\Sigma_{r_1}} X^{\otimes r_1}) \otimes \dots \otimes (a_s \otimes_{\Sigma_{r_s}} X^{\otimes r_s}) \subset \mathcal{P}(X)^{\otimes s}$$

where the first map permutes the factors and the second is the inclusion determined by $a_1 \otimes \dots \otimes a_s$. When k is a field of characteristic 0, linearity implies that these are the only possible natural transformations from $id^{\otimes r}$ to $\mathcal{P}(-)^{\otimes s}$, and hence that the map η is an isomorphism (see [27, App A, in particular (5.4)]—the case $s = 1$ is stated in [10, Prop 1.2.5], see also [11, Prop 2.3.12]). (Note that if we are not in characteristic 0, more maps are possible. For example the diagonal map $k \rightarrow k \otimes_{\Sigma_2} k$ is linear in characteristic 2, though not in characteristic 0.)

There is an isomorphism $\text{Hom}(id^{\otimes r}, \mathcal{P}(-)^{\otimes s}) \cong \text{Hom}((U_V)^{\otimes r}, (U_V)^{\otimes s})$, which takes a natural transformation $\theta = (\theta_X) \in \text{Hom}(id^{\otimes r}, \mathcal{P}(-)^{\otimes s})$ to the natural transformation given on a \mathcal{P} -algebra A by the map

$$U(A) = A^{\otimes r} \xrightarrow{\theta_{U(A)}} \mathcal{P}(A)^{\otimes s} \xrightarrow{\gamma_A^{\otimes s}} A^{\otimes s}$$

for $\gamma_A : \mathcal{P}(A) \rightarrow A$ the structure map of A , and with inverse taking a natural transformation $\eta = (\eta_A) \in \text{Hom}((U_V)^{\otimes r}, (U_V)^{\otimes s})$ to the natural transformation defined on a vector space X by

$$X^{\otimes r} \xrightarrow{i_X^{\otimes r}} \mathcal{P}(X)^{\otimes r} \xrightarrow{\eta_{\mathcal{P}(X)}} \mathcal{P}(X)^{\otimes s}$$

with $i_X : X \rightarrow id \otimes X \subset \mathcal{P}(X)$ the canonical inclusion. Now $\text{Hom}((U_V)^{\otimes r}, (U_V)^{\otimes s})$ is the completion of $\mathcal{E}_{\mathcal{P}}$ as defined above. The composition of these two isomorphisms gives

$$\mathcal{E}_{\mathcal{P}}(r, s) \xrightarrow{\cong} \text{Hom}(id^{\otimes r}, \mathcal{P}(-)^{\otimes s}) \cong \text{Hom}((U_V)^{\otimes r}, (U_V)^{\otimes s}) = \widehat{\mathcal{E}}_{\mathcal{P}}(r, s)$$

as claimed.

2.3. Example: the cap product. Given an algebra A (i.e. $A = \Phi(1)$ for $\Phi : \mathcal{A}ss \rightarrow \text{Ch}$ a symmetric monoidal functor), we can consider its endomorphism prop $\text{End}(A)$. It has morphisms from p to q defined as

$$\text{End}(A)(p, q) := \text{Hom}(A^{\otimes p}, A^{\otimes q})$$

where $A^{\otimes p}, A^{\otimes q}$ are considered as objects of Ch and Hom here denotes the chain complex of morphisms in that category. The algebra structure of A defines a symmetric monoidal functor $i : \mathcal{A}_\infty \rightarrow \text{End}(A)$ (factoring through $\mathcal{A}ss$), so $\text{End}(A)$ is a prop with \mathcal{A}_∞ -multiplication. The algebra A is by construction an $\text{End}(A)$ -algebra: $\text{End}(A)$ exactly encodes all the operations defined on A , and its associative algebra structure is the one obtained by pulling back its $\text{End}(A)$ -structure with i .

Evaluating on the $\text{End}(A)$ -algebra A , one sees that the restriction map

$$r : \text{Nat}_{\text{End}(A)} \longrightarrow \text{Nat}_{\text{End}(A)}^\otimes$$

is injective. Explicitly, $\text{Nat}_{\text{End}(A)}$ has morphism complexes

$$\text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right) \cong \prod_{j_1, \dots, j_{n_1} \geq 1} \bigoplus_{k_1, \dots, k_{n_2} \geq 1} \text{Hom}(A^{\otimes j_1 + \dots + j_{n_1} + m_1}, A^{\otimes k_1 + \dots + k_{n_2} + m_2})[k - j + n_1 - n_2]$$

with differential $d_A + d_H + d^H$ where d_H cyclically post-composes with multiplications and d^H cyclically precomposes with multiplications. (Recall from Example 1.4 that d^H is *not* the Hochschild cochain differential!)

Recall from [2, XI.6] the classical cap product in Hochschild homology:

$$\cap : C_p(A, A) \otimes C^q(A, A) \longrightarrow C_{p-q}(A, A).$$

Explicitly, given $a = a_0 \otimes \dots \otimes a_p \in C_p(A, A)$ and $D \in C^q(A, A) = \text{Hom}(A^{\otimes q}, A)$, it is given by the formula

$$(2) \quad a \cap D = (-1)^{(|a| - |a_0|)|D|} a_0 D(a_1, \dots, a_q) \otimes a_{q+1} \otimes \dots \otimes a_p$$

(see e.g. [29, Sec 2]). This operation is constructed using the $\text{End}(A)$ -algebra structure of A , and is thus part of the $\text{Nat}_{\text{End}(A)}$ -action:

Proposition 2.14. *There is an inclusion $F : C^*(A, A) \hookrightarrow \text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)$ satisfying that the composition*

$$C_*(A, A) \otimes C^*(A, A) \xrightarrow{id \otimes F} C_*(A, A) \otimes \text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) \longrightarrow C_*(A, A)$$

is the cap product.

Proof. Recall that $\text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) \cong \prod_{j \geq 1} \bigoplus_{k \geq 1} \text{Hom}(A^{\otimes j}, A^{\otimes k})[k - j]$. Given a Hochschild cochain $D \in \text{Hom}(A^q, A) = C^q(A, A)$, define $F(D)$ to have j th component the map $F(D)_j \in \text{Hom}(A^{\otimes j}, A^{\otimes j-q+1})$ defined by equation (2) above with $p = j - 1$, i.e. taking $a = a_0 \otimes \dots \otimes a_{j-1}$ to $a \cap D$. We need to check that $F(dD) = d(F(D))$. For simplicity, we assume A is concentrated in degree 0 and we leave the “non-structural” signs out. We have $(d(F(D)))_j = d_H(F(D)_j) + d^H(F(D)_{j-1})$. Now

$$(d_H(F(D)_j))(a) = d(a \cap D)$$

is the Hochschild differential applied to the cap product, and

$$(d^H(F(D)_{j-1}))(a) = -D \cap da.$$

As $(F(dD))(a) = dD \cap a$, the equation becomes

$$dD \cap a = d(a \cap D) - D \cap da$$

for each $a = a_0 \otimes \dots \otimes a_{j-1} \in A^{\otimes j}$ which is exactly saying that the cap product is a chain map. \square

Considering $C^*(A, A)$ as an algebra with the cup product, Nest and Tsygan define an action of $C_*(C^*(A, A), C^*(A, A))$ on $C_*(A, A)$ extending the cap product. This is further extended by Kontsevich-Soibelman in [21, Sec 11] to an action of $B(n, 1) \otimes C^*(A, A)$ on $C_*(A, A)$, where $B(n, 1)$ is a chain complex of certain configurations on a cylinder. In each case, the definition of the action, just as in the case of the cap product, gives maps

$$C_*(C^*(A, A), C^*(A, A)) \longrightarrow \text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)$$

and

$$B(n, 1) \otimes C^*(A, A) \longrightarrow \text{Nat}_{\text{End}(A)}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)$$

which are chain maps exactly because the actions are compatible with the differential.

3. THREE COMPUTATIONS OF THE FORMAL OPERATIONS

Let \mathcal{O} be the symmetric monoidal dg-category with objects the natural numbers and morphisms $\mathcal{O}(n, m)$ the chain complex of fat graphs with $n + m$ labeled leaves as described in Section 6. (See Figure 4 for an example of a morphism in \mathcal{O} .) Composition is defined by gluing leaves and the monoidal structure by disjoint union, just as for \mathcal{A}_∞ . In fact, the category \mathcal{A}_∞^+ identifies as a subcategory of forests in \mathcal{O} . The chain complex




FIGURE 4. Morphism in $\mathcal{O}(3, 4)$ (with the outputs marked as outgoing arrows)

$\mathcal{O}(n, m)$ computes the homology of the moduli space of Riemann cobordisms from n to m intervals, so that an \mathcal{O} -algebra, i.e. a strong symmetric monoidal functor $\Phi : \mathcal{O} \rightarrow \text{Ch}$, can be called an open topological conformal field theory (see [31, 30, 13, 4, 5] and [36, Thm 2.2]).

Let $H_0(\mathcal{O})$ denote the dg-category with the same objects as \mathcal{O} but with morphisms $H_0(\mathcal{O})(n, m) := H_0(\mathcal{O}(n, m))$. Explicitly, by the above, $H_0(\mathcal{O})(n, m)$ is the chain complex concentrated in degree 0 with one free generator for each topological type of cobordism from n to m intervals. A strong symmetric monoidal functor $\Phi : H_0(\mathcal{O}) \rightarrow \text{Ch}$, or $H_0(\mathcal{O})$ -algebra, is called an open topological quantum field theory, and such field theories are in 1-1 correspondence with symmetric Frobenius algebras (see [23, Cor 4.5]). Note that there is a quotient functor $\mathcal{O} \rightarrow H_0(\mathcal{O})$.

Including trees into all graphs defines a functor $i : \mathcal{A}_\infty^+ \rightarrow \mathcal{O}$, so \mathcal{O} and $H_0(\mathcal{O})$ are props with unital \mathcal{A}_∞ -multiplication and open field theories have a well-defined Hochschild complex. In the present section, we identify the chain complexes of formal operations $\overline{\text{Nat}}_{\mathcal{O}}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$, $\overline{\text{Nat}}_{H_0(\mathcal{O})}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$ and $\overline{\text{Nat}}_{\mathcal{A}_\infty^+}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$, up to homotopy, with well-known objects.

We start the section by analyzing the case of the category \mathcal{O} . The cases \mathcal{A}_∞^+ and $H_0(\mathcal{O})$ are then treated respectively as subcases and quotient of the case \mathcal{O} .

3.1. Open topological conformal field theories. Let \mathcal{O} be the open cobordism category described above and recall from Section 2 the chain complex $\overline{\text{Nat}}_{\mathcal{O}}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$ of formal operations on \mathcal{O} -algebras, the hom complex of the reduced Hochschild functors

$$\overline{\mathcal{C}}^{(n_i, m_i)} : \text{Fun}(\mathcal{O}, \text{Ch}) \rightarrow \text{Ch}.$$

We will relate this complex to a corresponding morphism complex in the open-closed cobordism category \mathcal{OC} , which we describe now.

For the purpose of the present paper, we could define \mathcal{OC} formally as a category with objects pairs of natural numbers $\begin{bmatrix} n \\ m \end{bmatrix}$, with $m, n \geq 0$, and morphism complexes

$$(3) \quad \mathcal{OC}\left(\begin{bmatrix} 0 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right) \cong \overline{\mathcal{C}}^{n_2}(\mathcal{O}(m_1, -))(m_2) = \bigoplus_{k_1, \dots, k_{n_2}} \mathcal{O}(m_1, k + m_2)/_U \otimes L_{k_1} \otimes \dots \otimes L_{k_{n_2}},$$

the reduced iterated Hochschild complex of $\mathcal{O}(m_1, -) : \mathcal{O} \rightarrow \text{Ch}$ at m_2 , and with $\mathcal{OC}\left(\begin{bmatrix} n_1 \\ m_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$ for $n_1 > 0$ a certain subcomplex of $\mathcal{OC}\left(\begin{bmatrix} 0 \\ m_1 + n_1 \end{bmatrix}, \begin{bmatrix} n_2 \\ m_2 \end{bmatrix}\right)$, together with a certain composition law. (Here $k = k_1 + \dots + k_{n_1}$ and L_{k_i} is the free module on one

generator l_{k_i} in degree $k_i - 1$ of Section 1.) The fact that it can be described this way is a major ingredient in our identification of the natural operations, but it would be difficult to describe which subcomplex $\mathcal{OC}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$ really is, or define the composition, without introducing open-closed cobordisms and black and white graphs, which we do now. At the same time, we will give geometric meaning to the category \mathcal{OC} .

An open-closed cobordism is a surface S whose boundary has an incoming and outgoing part: $\partial S \supset \partial_{in} S \sqcup \partial_{out} S$, where $\partial_{in} S$ and $\partial_{out} S$ are disjoint unions of labeled circles and closed intervals in ∂S , with S then considered as a cobordism from $\partial_{in} S$ to $\partial_{out} S$. The part of the boundary which is neither incoming nor outgoing is called *free*: $\partial_{free} S = \partial S \setminus (\partial_{in} S \cup \partial_{out} S)$. The open-closed cobordism category \mathcal{OC} considered here has objects pairs of natural numbers $\begin{smallmatrix} n \\ m \end{smallmatrix}$, thought of as a union of m intervals and n circles, and morphisms $\mathcal{OC}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$ a chain complex whose homology is that of the moduli space of Riemann cobordisms from m_1 intervals and n_1 circles to m_2 intervals and n_2 circles satisfying that each component has non-empty free or incoming boundary.

To describe \mathcal{OC} explicitly, we use the language of black and white graphs. A *black and white graph* is a fat graph whose set of vertices is given as $V = V_b \sqcup V_w$, with V_b the set of *black* vertices and V_w the set of *white* vertices. Black vertices are assumed to be at least trivalent whereas white vertices are allowed to be any valence at least 1. In addition, white vertices are labeled, and come equipped with the data of a start half-edge at the vertex. The graph l_n from Section 1 is a black and white graph. Figure 5(a) gives a more generic example. Figure 5(b) shows how to associate a surface with boundaries to a black and white graph, fattening the graph and inserting boundary circles where there were white vertices. A *boundary cycle* in the graph is a sequence of half-edges in the graph corresponding to a boundary component of that surface not coming from a white vertex.




FIGURE 5. Black and white graph and associated cobordism.

Black and white graphs form a chain complex just like fat graphs, defining the degree of a graph as $\sum_{v \in V_b} (|v| - 3) + \sum_{v \in V_w} (|v| - 1)$, and with the differential given as sums of all possible blow-ups, a white vertex blowing up as a pair of a white and a black vertex. In fact, one can think of a black and white graph as made out of one copy of the graph $l_{|v|}$ of Section 1 for each white vertex v , attached along a fat graph, and the differential at such a white vertex v is that of $l_{|v|}$ —This translates to the formal description of \mathcal{OC} given in equation (3) above. We have also mentioned this correspondence earlier in Remark 1.1 when describing the functor \mathcal{L} used to define the Hochschild complex. We refer to [36, Sec 2.3, 2.5] for more details about the chain complex of black and white graphs.

Now $\mathcal{OC}(\begin{smallmatrix} n_1 \\ m_1 \end{smallmatrix}, \begin{smallmatrix} n_2 \\ m_2 \end{smallmatrix})$ is a chain complex generated by black and white graphs with n_2 white vertices modeling the n_2 outgoing closed boundary components, and $n_1 + m_1 + m_2$ labeled leaves such that the first n_1 leaves each are sole leaves in their boundary cycle, representing the n_1 closed incoming, m_1 open incoming and m_2 open outgoing boundaries. Composition is obtained by summing over all ways of removing the white vertices of the first graph (its outgoing closed boundaries) and attaching their leaves to the corresponding incoming boundary cycle of the second graph, in a way that respects

the cyclic ordering and preserves the grading, and finally attaching in pairs the leaves modeling open boundaries as in \mathcal{O} . We refer to [36, Sec 2.8] for further details about the category \mathcal{OC} , including a precise definition of the composition. It follows from [4, Prop 6.1.3] that this chain complex computes the homology of the moduli space of Riemann cobordisms from m_1 intervals and n_1 circles to m_2 intervals and n_2 circles satisfying that each component has at least an incoming or a free boundary (see [36, Thm 5.4]). The fact that composition in \mathcal{OC} corresponds to gluing of surfaces on the moduli space is shown in [7].

Cutting around the white vertices yields the isomorphism

$$\mathcal{OC}([m_1^0], [m_2^{n_2}]) \cong \overline{\mathcal{C}}^{n_2}(\mathcal{O}(m_1, -))(m_2) = \bigoplus_{k_1, \dots, k_{n_2}} \mathcal{O}(m_1, k + m_2)/U \otimes L_{k_1} \otimes \cdots \otimes L_{k_{n_2}}$$

mentioned above. (This is the content of [36, Lem 6.1].) In the language of [36], this means that \mathcal{OC} is an extension of the Hochschild core category of \mathcal{O} . Corollary 5.12 of [36] can then be rephrased as saying that \mathcal{OC} acts naturally on the functors

$$\overline{\mathcal{C}}^n(-)(m) : \text{Fun}(\mathcal{O}, \text{Ch}) \longrightarrow \text{Ch}$$

i.e. that there is a functor

$$J_{\mathcal{O}} : \mathcal{OC} \rightarrow \overline{\text{Nat}}_{\mathcal{O}}.$$

To describe this functor explicitly on morphism complexes, we first rewrite its target using the identifications from Theorem 2.1 and [36, Lem. 6.1]:

$$\overline{\text{Nat}}_{\mathcal{O}}([m_1^{n_1}], [m_2^{n_2}]) \cong \overline{\mathcal{D}}^{n_1} \overline{\mathcal{C}}^{n_2} \mathcal{O}(-, -)(m_1, m_2) \cong \overline{\mathcal{D}}^{n_1}(\mathcal{OC}([m_1^0], [m_2^{n_2}]))(m_1)$$

which exhibits $\overline{\text{Nat}}_{\mathcal{O}}([m_1^{n_1}], [m_2^{n_2}])$ as a subcomplex of the product $\prod_{\underline{j}} \mathcal{OC}([m_1^0 + j], [m_2^{n_2}])[n_1 - j]$ over $\underline{j} = (j_1, \dots, j_{n_1})$ and with $j = j_1 + \cdots + j_{n_1}$ as before. Then the map

$$J_{\mathcal{O}} : \mathcal{OC}([m_1^{n_1}], [m_2^{n_2}]) \longrightarrow \overline{\text{Nat}}_{\mathcal{O}}([m_1^{n_1}], [m_2^{n_2}]) \subset \prod_{\underline{j}} \mathcal{OC}([m_1^0 + j], [m_2^{n_2}])[n_1 - j]$$

can be explicitly identified as the map that takes a graph G to the sequence $\{G \circ (l_{\underline{j}} + id_{m_1})\}_{\underline{j}}$, where $l_{\underline{j}} = l_{j_1} \otimes \cdots \otimes l_{j_{n_1}} \in L_{j_1} \otimes \cdots \otimes L_{j_{n_1}}$ corresponds in $\mathcal{OC}([m_1^0], [m_2^{n_1}])$ to the black and white graph defined by the disjoint union of the graphs l_{j_i} , with \circ the composition in \mathcal{OC} . More explicitly, the j th component of the image of G is the sum of all graphs that can be obtained from G by gluing $l_{j_i} - 1$ labeled leaves cyclically at the vertices of the boundary cycle of G corresponding to its i th incoming closed boundary for each $i = 1, \dots, n_1$. One can check that this map has image in the reduced coHochschild complex.

Note that the map $J_{\mathcal{O}}$ is injective on each morphism complex. A splitting can be constructed using the map $\overline{\text{Nat}}_{\mathcal{O}}([m_1^{n_1}], [m_2^{n_2}]) \rightarrow \mathcal{OC}([m_1^0 + n_1], [m_2^{n_2}])$ associating to a sequence $\{G_{\underline{j}}\}_{\underline{j}} \in \prod_{\underline{j}} \mathcal{OC}([m_1^0 + j], [m_2^{n_2}])[n_1 - j]$ its \underline{j} th component for $\underline{j} = (1, \dots, 1)$. The categories are however far from being isomorphic. In particular, the morphisms complexes in $\overline{\text{Nat}}_{\mathcal{O}}$, unlike in \mathcal{OC} , are not positively graded. We will however show the following result:

Theorem 3.1. *The functor $J_{\mathcal{O}} : \mathcal{OC} \xrightarrow{\sim} \overline{\text{Nat}}_{\mathcal{O}}$ is a quasi-isomorphism of categories, which is split injective on each morphism complex.*

The main ingredient in the proof is a description of the complex $\overline{\mathcal{D}}^{n_1}(\mathcal{OC}([m_1^0], [m_2^{n_2}]))(m_1)$ in terms of a cosimplicial set of partitions, which we define first.

Let X be an oriented 1-manifold without boundary (i.e. a collection of circles and open intervals). Let $K(q, X) = \pi_0 \text{Conf}(q, X)$ denote the components of the configuration space of q ordered points in X . These can be thought of as partitions of $\{1, \dots, q\}$ which are ordered on the interval components of X and cyclically ordered on the circle components.

Lemma 3.2. *For each oriented 1-manifold X without boundary, the sets $\{\mathbb{K}(q+1, X)\}_{q \geq 0}$ form a cosimplicial set.*

Proof. Let $(p_0, \dots, p_q) \in X^{q+1}$ represent an element in $\mathbb{K}(q+1, X)$. Define

$$d^i : \mathbb{K}(q+1, X) \rightarrow \mathbb{K}(q+2, X)$$

by adding a copy \tilde{p}_i of p_i to the right of p_i for $0 \leq i \leq q$, and by adding a copy \hat{p}_0 of p_0 to the left of p_0 for $i = q+1$, to get a new configuration represented by $(p_0, \dots, p_i, \tilde{p}_i, \dots, p_q)$ (resp. $(p_0, \dots, p_q, \hat{p}_0)$). Define also

$$s^i : \mathbb{K}(q+1, X) \rightarrow \mathbb{K}(q, X)$$

forgetting p_{i+1} , $0 \leq i \leq q-1$, and relabeling respecting the order.

We are left to check the cosimplicial identities:

$$\begin{aligned} d^j d^i &= d^i d^{j-1} & i < j \\ s^j d^i &= d^i s^{j-1} & i < j \\ s^j d^j &= 1 = s^j d^{j+1} \\ s^j d^i &= d^{i-1} s^j & i > j+1 \\ s^j s^i &= s^i s^{j+1} & i \leq j \end{aligned}$$

For the first identity, if $j \leq q+1$, both compositions repeat p_i and p_{j-1} to their right and if $j = q+2$, both compositions repeat p_i and p_0 , one to the right, the other one to the left, or they repeat twice p_0 to its left if $i = q+1$. For the second, both compositions double p_i to its right and forget p_j . For the third, the first composition adds a copy of p_j and forgets the added point, while the second composition adds a copy of p_{j+1} and forgets the original p_{j+1} if $j < q$, or, when $j = q$, it adds a copy of p_0 to its right and forgets it. For the fourth equality, both compositions forget p_{j+1} and add p_i to its right (if $i \leq q$) or p_0 to its left (if $i = q+1$). Finally in the last one, both compositions forget p_{i+1} and p_{j+2} . \square

There is a chain complex $\bigoplus_q \mathbb{Z} \mathbb{K}(q+1, X)$ associated to the cosimplicial set $\mathbb{K}(\bullet+1, X)$, with $\mathbb{Z} \mathbb{K}(q+1, X)$ in degree $-q$, and with differential the alternating sum of the coface maps

$$d_K = \sum (-1)^i d^i : \mathbb{Z} \mathbb{K}(q, X) \rightarrow \mathbb{Z} \mathbb{K}(q+1, X)$$

raising the degree by one. The appendix Section 5 studies the properties of such chain complexes. In particular, Proposition 5.4 says that such a chain complex has homology concentrated in degree 0.

Let $\mathbb{K}_c(q, X)$ denote the subset of $\mathbb{K}(q, X)$ of *complete configurations*, those with at least one point in each circle component of X . Note that the coface maps d^i restrict to the subsets $\mathbb{K}_c(q, X)$. In fact, for $x \in \mathbb{K}(q, X)$ with $d_K(x) \neq 0$, we have $d_K(x) \in \mathbb{Z} \mathbb{K}_c(q+1, X)$ if and only if $x \in \mathbb{K}_c(q, X)$, so that the subcomplex $(\mathbb{Z} \mathbb{K}_c(*, X), d_K)$ splits off as a union of components of the chain complex $(\mathbb{Z} \mathbb{K}(*, X), d_K)$. We reinterpret $(\mathbb{Z} \mathbb{K}_c(*, X), d_K)$ as the coHochschild complex of a functor $\mathbb{Z} \mathbb{K}_c(-, X)$ as follows: with $\mathcal{A}ss$ denoting the prop of (non-unital) associative algebras, define

$$\mathbb{Z} \mathbb{K}_c(-, X) : (\mathcal{A}ss)^{op} \longrightarrow \text{Ch}$$

on objects by $\mathbb{Z} \mathbb{K}_c(-, X)(n) = \mathbb{Z} \mathbb{K}_c(n, X)$ concentrated in degree 0, and on morphisms by doubling and relabeling points in the configurations according to the morphisms in $\mathcal{A}ss$. As $\mathbb{Z} \mathbb{K}_c(-, X)(n)$ is concentrated in degree 0 for each n , we have that its coHochschild complex is really a direct sum complex, with $\mathbb{Z} \mathbb{K}_c(n, X)$ in degree $1-n$. In fact we have the following:

Lemma 3.3. *For any oriented 1-manifold without boundary X , we have an isomorphism $(\mathbb{Z} \mathbb{K}_c(*, X), d_K) \cong D(\mathbb{Z} \mathbb{K}_c(-, X))(0)$ between the chain complex of the cosimplicial set of complete configurations in X , and the coHochschild complex of the associated functor.*

For each n_1, n_2, m_1, m_2 , we consider the functor

$$\mathcal{OC}([m_1+q], [m_2]) : \mathcal{O}^{op} \longrightarrow \text{Ch}$$

taking the value $\mathcal{OC}([m_1+q], [m_2])$ on the object q in \mathcal{O} , and defined on morphisms by precomposition in \mathcal{OC} via the inclusion of \mathcal{O} in $\mathcal{OC}([m_1+q], [m_2])$ tensoring with the identity on $[m_1]$.

We give now a model for the restriction of $\mathcal{OC}([m_1+q], [m_2])$ to \mathcal{A}_∞^{op} which splits out the configuration information defined by the incoming and outgoing boundary components of a cobordism: Let

$$(\mathcal{OC} \otimes \text{K})([m_1], [m_2]) : (\mathcal{Ass})^{op} \longrightarrow \text{Ch}$$

be defined on the object q in \mathcal{Ass} by

$$(\mathcal{OC} \otimes \text{K})([m_1], [m_2])(q) = \bigoplus_{b=0}^q \bigoplus_S \mathcal{OC}_S([m_1+b], [m_2]) \otimes \mathbb{Z} \text{K}_c(q, \coprod_b S^1 \cup \partial_I S)$$

where S runs over the components of $\mathcal{OC}([m_1+b], [m_2])$, and \mathcal{OC}_S denotes the corresponding component, and $\partial_I S$ denotes the interval components of the free boundary of S . The differential is the differential in \mathcal{OC} . On morphisms, \mathcal{Ass} acts via its action on the second factor.

We can pull back the above two functors via $i : \mathcal{A}_\infty \rightarrow \mathcal{O}$ and $j : \mathcal{A}_\infty \rightarrow \mathcal{Ass}$ respectively to functors defined on $(\mathcal{A}_\infty)^{op}$.

Lemma 3.4. *There is a quasi-isomorphism of functors*

$$i^* \mathcal{OC}([m_1+q], [m_2]) \xrightarrow{\cong} j^*(\mathcal{OC} \otimes \text{K})([m_1], [m_2]) : (\mathcal{A}_\infty)^{op} \longrightarrow \text{Ch}.$$

Proof. We first define for each q a map

$$\beta_q : \mathcal{OC}([m_1+q], [m_2]) \longrightarrow \bigoplus_{b=0}^q \bigoplus_S \mathcal{OC}_S([m_1+b], [m_2]) \otimes \text{K}_c(q, \coprod_b S^1 \cup \partial_I S).$$

Recall that a closed incoming boundary component is modeled in \mathcal{OC} by a leaf alone in its boundary cycle in the graph. A graph $G \in \mathcal{OC}([m_1+q], [m_2])$ is a graph with n_2 white vertices, q leaves p_1, \dots, p_q and $n_1 + m_1 + m_2$ leaves $\lambda_1, \dots, \lambda_{n_1+m_1+m_2}$, with the first n_1 leaves alone in their boundary cycle. Let b be the number of boundary cycle of G having leaves only of type p_i . Define $\beta_q(G) = G' \otimes P$, where G' is obtained from G by forgetting the leaves p_1, \dots, p_q (and their attaching trees) except for p_{i_1}, \dots, p_{i_b} , the first occurring leaf of that type in each boundary cycle of G with only leaves labeled by p_i 's, provided that G and G' have the same degree, i.e. provided that the forgotten leaves were attached at trivalent black vertices or univalent white vertices. Otherwise we set $\beta_q(G) = 0$. Let S be the topological cobordism associated to G' . The second component, P , remembers the configuration defined by the leaves p_i in ∂S . This configuration is supported in $\coprod_b S^1 \cup \partial_I S$. (See Figure 6 for an example.)




FIGURE 6. Graph G with $n_1 = 1$, $n_2 = 2$, $m_1 = 1$, $m_2 = 2$ and $q = 6$, and its image $G' \otimes P$ under β_6 , with S the underlying cobordism type of G' , where the thicker lines show the support of P .

We check that β_q is a chain map, i.e. that it commutes with the differential in \mathcal{OC} . If $\beta_q(G) = G' \otimes P$ with $G' \neq 0$, then $\beta_q(d_{\mathcal{OC}}G) = d_{\mathcal{OC}}G' \otimes P$ as the forgotten leaves

where attached at trivalent black vertices or univalent white vertices in that case. On the other hand, if $\beta_q(G) = 0$, then each component of $d_{\mathcal{OC}}G$ will also be mapped to 0 by β_q , unless G has a single valence 4 black vertex or valence 2 white vertex with a p_i forgotten by β_q . In this case, there are two terms in $d_{\mathcal{OC}}G$ on which β_q does not vanish, but their images will be isomorphic and with opposite orientations. Hence each β_q is a chain map.

Furthermore, the β_q 's assemble to a natural transformation of functors: for $f \in (\mathcal{A}_\infty)^{op}$, $\beta(f(G)) = j(f)\beta(G)$ as the image of a graph G under positive degree morphisms of \mathcal{A}_∞ are killed by β , and the action of degree 0 morphisms on graphs corresponds exactly to the action on K_c via the projection $j : \mathcal{A}_\infty \rightarrow \mathcal{Ass}$.

We are left to show that each β_q is a quasi-isomorphism. Note first that the above map is an isomorphism on sets of components as there is exactly one topological type of cobordism on the left hand side corresponding to each pair (S, P) on the right hand side.

Finally, the homology groups on the left hand side only depend on b , not on the particular partition in $K_c(\bullet + 1, \partial_I S \cup \partial_{[b]} S)$ as the mapping class group of a surface fixing any number of points in a boundary component is isomorphic to the mapping class group fixing the whole boundary. \square

We are now ready to prove our second lemma, which, iterated n_1 times, will prove the theorem.

Lemma 3.5. *There is a quasi-isomorphism of functors*

$$\mathcal{OC}([_{-}^{n_1+1}], [_{m_2}^{n_2}]) \xrightarrow{\simeq} D(\mathcal{OC}([_{-}^{n_1}], [_{m_2}^{n_2}]))(-) : \mathcal{O}^{op} \rightarrow \text{Ch}$$

defined at the object m_1 by taking a graph G to $\{G \circ (l_j + id_{m_1})\}_{j \geq 1}$, using the identification $D(\mathcal{OC}([_{-}^{n_1}], [_{m_2}^{n_2}]))(m_1) \cong \prod_{j \geq 1} \mathcal{OC}([_{m_1+j}^{n_1}], [_{m_2}^{n_2}])[1-j]$.

(The map does actually have image in the quasi-isomorphic reduced subcomplex.)

Proof. Note first that the map is indeed a natural transformation as \mathcal{O} acts by composition on boundaries that are not affected by the glued l_j 's.

We are left to check that it induces isomorphisms in homology pointwise. We have $D(\mathcal{OC}([_{-}^{n_1}], [_{m_2}^{n_2}]))(m_1) \cong D(\mathcal{OC}([_{m_1+}^{n_1}], [_{m_2}^{n_2}]))(0)$. By Lemma 3.4 and the homotopy invariance of the coHochschild construction (Proposition 1.5), we thus get

$$D(\mathcal{OC}([_{-}^{n_1}], [_{m_2}^{n_2}]))(m_1) \simeq D((\mathcal{OC} \otimes K)([_{m_1}^{n_1}], [_{m_2}^{n_2}]))(0).$$

Recall from Section 1.1 the filtration

$$F^s = \prod_{q \geq s} \bigoplus_{b=0}^q \bigoplus_S \mathcal{OC}_S([_{m_1}^{n_1+b}], [_{m_2}^{n_2}]) \otimes \mathbb{Z} K_c(q, \coprod_b S^1 \cup \partial_I S)$$

which is exhaustive and complete. Its E^1 -term has

$$E_{-p,q}^1 = \bigoplus_{b=0}^{p+1} \bigoplus_S H_q(\mathcal{OC}_S([_{m_1}^{n_1+b}], [_{m_2}^{n_2}])) \otimes \mathbb{Z} K_c(p+1, \coprod_b S^1 \cup \partial_I S)$$

and the d^1 -differential is the coHochschild differential, which, by Lemma 3.3, can be identified with the differential of the chain complex associated to the cosimplicial set $K(\bullet, \coprod_b S^1 \cup \partial_I S)$, restricted to the split subcomplex $\mathbb{Z} K_c$.

By Proposition 5.4, the homology of this complex is concentrated in degree 0, with a summand for each element x in $K_c(1, \coprod_b S^1 \cup \partial_I S)$ satisfying that $d^0 x = d^1 x$. Such an x is a configuration of a single element p_0 , and d^0, d^1 double it as (p_0, p_1) and (p_1, p_0) . These two configurations are in the same component exactly when p_0 is alone on its

boundary component. There is exactly one such configuration in $K_c(1, \coprod_b S^1 \cup \partial_I S)$ when $b = 1$ and none if $b \neq 1$. Hence the E^2 -term is given as $E_{-p,q}^2 = 0$ for $p > 0$ and

$$E_{0,q}^2 = H_q(\mathcal{OC}([{}_{m_1}^{n_1+1}], [{}_{m_2}^{n_2}])).$$

Now using the Eilenberg-Moore comparison theorem ([37, Thm 5.5.11]) for the trivial filtration of $H_*(\mathcal{OC}([{}_{m_1}^{n_1+1}], [{}_{m_2}^{n_2}]))$ given by $F^1 = H_*(\mathcal{OC}([{}_{m_1}^{n_1+1}], [{}_{m_2}^{n_2}])), F^s = 0$ for $s > 1$, the map in the statement of the corollary gives an isomorphism on the E^2 -pages, and hence an isomorphism $H_*(\mathcal{OC}([{}_{m_1}^{n_1+1}], [{}_{m_2}^{n_2}])) \xrightarrow{\cong} H_*(D((\mathcal{OC} \otimes K)([{}_{-}^{n_1}], [{}_{m_2}^{n_2}]))(m_1))$. \square

Proof of Theorem 3.1. Applying Lemma 3.5 n_1 times together with the homotopy invariance of D (Proposition 1.5), we get a sequence of quasi-isomorphisms

$$\mathcal{OC}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}]) \xrightarrow{\cong} D(\mathcal{OC}([{}_{-}^{n_1-1}], [{}_{m_2}^{n_2}]))(m_1) \xrightarrow{\cong} \dots \xrightarrow{\cong} D^{n_1}(\mathcal{OC}([{}_{-}^0], [{}_{m_2}^{n_2}]))(m_1)$$

whose composition is the map given in the statement of the theorem. As that map has image in the subcomplex $\overline{\text{Nat}}_{\mathcal{O}}(\mathcal{OC}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}])) = \overline{D}^{n_1}(\mathcal{OC}([{}_{-}^0], [{}_{m_2}^{n_2}]))(m_1)$, which is quasi-isomorphic to $D^{n_1}(\mathcal{OC}([{}_{-}^0], [{}_{m_2}^{n_2}]))(m_1)$ by Proposition 1.7, we get the desired quasi-isomorphism. \square

Remark 3.6. There is a space-level version of the above algebraic phenomenon. Given a surface S with $n + m$ boundary components $\partial_n S \cup \partial_m S$, with $m \geq 1$, there is a cosimplicial space $\mathcal{K}^\bullet(S, n)$ with

$$\mathcal{K}^q(S, n) := \coprod_{p_0, \dots, p_k \in \partial_m S} B\Gamma(S; \partial_n S \cup \{p_0, \dots, p_k\})$$

where $\Gamma(S; \partial_n S \cup \{p_0, \dots, p_k\})$ denotes the mapping class group of S fixing the first n boundary components of S pointwise as well as fixing each of the marked points p_0, \dots, p_k lying in the last m boundary components of S . The coface and codegeneracy maps are defined as for the cosimplicial set $K(\bullet, X)$ by doubling and forgetting marked points. One can then check that $\text{Tot}(\mathcal{K}^\bullet(S, n)) \cong B\Gamma(S; \partial_{n+1} S)$.

3.2. Unital \mathcal{A}_∞ -algebras. We want to restrict the argument of the previous section for the open cobordism category \mathcal{O} to its subcategory \mathcal{A}_∞^+ . Recall from [36, Prop 6.12] that

$$\overline{C}^{n_2}(\mathcal{A}_\infty^+(m_1, -)(m_2)) \cong \bigoplus_A \mathcal{OC}_A([{}_{m_1}^0], [{}_{m_2}^{n_2}])$$

is a sum of components of \mathcal{OC} , where A runs over the cobordisms which are a union of m_2 discs, each with exactly one open outgoing boundary, and n_2 annuli, each with precisely one closed outgoing boundary, and with a total of m_1 incoming open boundaries distributed on the free boundaries of the discs and annuli. In particular, we can now identify

$$\overline{\text{Nat}}_{\mathcal{A}_\infty^+}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}]) \cong \overline{D}^{n_1}(\bigoplus_A \mathcal{OC}_A([{}_{-}^0], [{}_{m_2}^{n_2}]))(m_1)$$

which exhibits $\overline{\text{Nat}}_{\mathcal{O}}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}])$ as a subcomplex of $\prod_{j_1, \dots, j_{n_1}} \bigoplus_A \mathcal{OC}_A([{}_{m_1+j}^0], [{}_{m_2}^{n_2}])[n_1 - j]$.

Let Ann denote the subcategory of \mathcal{OC} with the same objects, with

$$\text{Ann}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}]) \cong \bigoplus_{A'} \mathcal{OC}_{A'}([{}_{m_1}^{n_1}], [{}_{m_2}^{n_2}])$$

where A' runs over cobordisms which are a union of m_2 discs and $n_2 - n_1$ annuli as before, union with n_1 annuli each with one incoming closed and one outgoing closed boundary components. This chain complex can thus also be identified with $\bigoplus_A \mathcal{OC}_A([{}_{m_1}^0], [{}_{m_2}^{n_2-n_1}]) \otimes$

$\mathcal{OC}_{S^1 \times I}(\begin{smallmatrix} [1] \\ [0] \end{smallmatrix}, \begin{smallmatrix} [1] \\ [0] \end{smallmatrix})^{\otimes n_1}$, with A as above. Restriction of the functor $J_{\mathcal{O}}$ of the previous section of Ann yields a commutative diagram:

$$\begin{array}{ccc} \text{Ann}(\begin{smallmatrix} [n_1] \\ [m_1] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix}) & \xrightarrow{J_{\mathcal{A}_{\infty}^+}} & \overline{\text{Nat}}_{\mathcal{A}_{\infty}^+}(\begin{smallmatrix} [n_1] \\ [m_1] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix}) \hookrightarrow \prod_{\underline{j}} \bigoplus_A \mathcal{OC}_A(\begin{smallmatrix} 0 \\ [m_1+j] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix})[n_1 - j] \\ \downarrow & & \downarrow \\ \mathcal{O}(\begin{smallmatrix} [n_1] \\ [m_1] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix}) & \xrightarrow{J_{\mathcal{O}}} & \overline{\text{Nat}}_{\mathcal{O}}(\begin{smallmatrix} [n_1] \\ [m_1] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix}) \hookrightarrow \prod_{\underline{j}} \mathcal{OC}(\begin{smallmatrix} 0 \\ [m_1+j] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix})[n_1 - j] \end{array}$$

where $J_{\mathcal{A}_{\infty}^+}$, the restriction of $J_{\mathcal{O}}$ to Ann , takes a graph G to the sequence $\{G \circ (l_{\underline{j}} + id_{m_1})\}_j$, where $G \circ (l_{\underline{j}} + id_{m_1})$ is obtained from G by replacing its i th incoming closed boundary by j_i cyclically ordered incoming open boundaries for each $0 \leq i \leq n_1$.

The proof of Theorem 3.1 restricts to the annuli components arising above to prove that $J_{\mathcal{A}_{\infty}^+}$ is also a quasi-isomorphism:

Theorem 3.7. *The functor $J_{\mathcal{A}_{\infty}^+} : \text{Ann} \rightarrow \overline{\text{Nat}}_{\mathcal{A}_{\infty}^+}$ is a quasi-isomorphism of categories.*

Recall from [36, Prop 6.14] that $H_*(\text{Ann})$ is generated as a symmetric monoidal category by the unit and multiplication of the algebra, the Δ -operator representing Connes-Rinehart's B operator on the Hochschild complex of an associative algebra, and the inclusion of the algebra in its Hochschild complex.

3.3. Strict Frobenius algebras. Let $H_0(\mathcal{O})$ be the dg-category obtained from \mathcal{O} by taking its 0th homology, as described at the beginning of the section, whose algebras are symmetric Frobenius algebras. We will describe the formal operations on $H_0(\mathcal{O})$ -algebras in terms of Sullivan diagrams.

Following [36], we define here *Sullivan diagrams* as equivalence classes of black and white graphs having only trivalent black vertices, where the equivalence relation is generated by the boundary of graphs with a single valence 4 black vertex. By [36, Thm 2.6], this is equivalent to the more classical definition, which, loosely speaking, defines Sullivan diagrams as equivalence classes of fat graphs build from a number of circles by attaching chords, where the chords are thought of as having length 0. (Figure 8 shows examples of a few Sullivan diagrams both represented as black and white graphs and as ‘‘classical’’ Sullivan diagrams. The white vertices in the black and white graphs description correspond to the circles in the classical picture.) Sullivan diagrams model the unimodular harmonic compactification of moduli space [8].

The category \mathcal{SD} of Sullivan diagram can then be directly defined as the quotient category of the open-closed category \mathcal{OC} by the graphs with higher valence black vertices, and by the boundaries of such. (See [36, Sec 2.10] for a less concise description of that category.)

Lemma 6.6 of [36] shows that, cutting around the white vertices gives an isomorphism

$$\mathcal{SD}(\begin{smallmatrix} 0 \\ [m_1] \end{smallmatrix}, \begin{smallmatrix} [n_2] \\ [m_2] \end{smallmatrix}) \cong \overline{\mathcal{C}}^{n_2}(H_0(\mathcal{O})(m_1, -))(m_2)$$

which says, in the language of [36], that \mathcal{SD} is an extension of the Hochschild core category of $H_0(\mathcal{O})$. Hence by [36, Cor 5.12], \mathcal{SD} acts naturally on the functors $\overline{\mathcal{C}}^{(n,m)} : \text{Fun}(H_0(\mathcal{O}), \text{Ch}) \rightarrow \text{Ch}$, that is there is a functor

$$J_{H_0} : \mathcal{SD} \longrightarrow \overline{\text{Nat}}_{H_0(\mathcal{O})}$$

which is an inclusion on morphism complexes just as in the case of the prop \mathcal{O} . We prove here that this functor is a quasi-isomorphism:

Theorem 3.8. *The functor $J_{H_0} : \mathcal{SD} \xrightarrow{\sim} \overline{\text{Nat}}_{H_0(\mathcal{O})}$ is a quasi-isomorphism of categories, which is split-injective on each morphism complex.*

We will prove this theorem by adapting the proof of Theorem 3.1 to the present case. We first need the analogue of Lemma 3.4.

Note that the components of the morphism complexes of Sullivan diagrams are in one to one correspondence with those of \mathcal{OC} as the equivalence relation defining Sullivan diagrams respects the components.

Just as in the case of the category \mathcal{OC} , let

$$(\mathcal{SD} \otimes K)([m_1], [m_2]) : (\mathcal{Ass})^{op} \longrightarrow \text{Ch}$$

be defined on objects by

$$(\mathcal{SD} \otimes K)([m_1], [m_2])(q) = \bigoplus_{b=0}^q \bigoplus_S \mathcal{SD}_S([m_1+b], [m_2]) \otimes \mathbb{Z} K_c(q, \coprod_b S^1 \cup \partial_I S)$$

where S runs over the components of $\mathcal{SD}([m_1+b], [m_2])$, and $\partial_I S$ denotes the interval components of the free boundary of S . The differential is the differential in \mathcal{SD} . On morphisms, \mathcal{Ass} acts via its action on the second factor.

Lemma 3.9. *There is a quasi-isomorphism of functors*

$$i^* \mathcal{SD}([m_1+-, [m_2]) \simeq j^* (\mathcal{SD} \otimes K)([m_1], [m_2]) : (\mathcal{A}_\infty)^{op} \longrightarrow \text{Ch}.$$

Proof. The map

$$\beta_q : \mathcal{SD}([m_1+q], [m_2]) \longrightarrow \bigoplus_{b=0}^q \bigoplus_S \mathcal{SD}_S([m_1+b], [m_2]) \otimes K_c(q, \coprod_b S^1 \cup \partial_I S).$$

is defined as in the case of \mathcal{OC} by taking a Sullivan diagram G with n_2 white vertices and $q+1$ leaves p_0, \dots, p_q and $m_1+m_2+n_1$ leaves $\lambda_1, \dots, \lambda_{m_1+m_2+n_1}$ to $G' \otimes P$ where G' is obtained from G by forgetting the leaves p_0, \dots, p_q except for p_{i_1}, \dots, p_{i_b} , the first occurring leaf in each boundary cycle of G with only leaves labeled by p_i 's, provided that G and G' have the same degree. Otherwise we set $G' = 0$. The second component, P , remembers the configuration defined by the leaves p_i in ∂S . The map β_q respects the equivalence relation defining Sullivan diagrams and is a chain map as it is a quotient of the same map for \mathcal{OC} . We are left to check that it is a quasi-isomorphism.

In the present case, there is a right inverse γ_q defined by taking $G \otimes P$ to the graph obtained from G by adding the forgotten leaves recorded in P as a trivalent tree, in the ordering prescribed by P (as in Figure 7). This is a natural transformation as the




FIGURE 7. The maps β_q and γ_q for Sullivan diagrams

coHochschild and cosimplicial differentials agree under this map using the equivalence relation of Sullivan diagrams.

We have $\beta_q \circ \gamma_q = id$, so we are left to show that $\gamma_q \circ \beta_q \simeq id$. The failure of $\gamma_q \circ \beta_q$ to be the identity is on graphs with leaves that are “at or past some white vertices” of their place after applying $\gamma_q \circ \beta_q$. The chain homotopy is defined by summing over the graphs with each leaf, one at a time, at the white vertex which has to be past, for each white vertex. This defines a degree 1 map $s : \mathcal{SD}([m_1+q], [m_2]) \rightarrow \mathcal{SD}([m_1+q], [m_2])$ and we need to check that $sd + ds = id - \gamma_q \circ \beta_q$. In fact, it is enough to check that we can pass leaves one at a time: (we omit the sign calculation) for a graph G with a leaf λ_1 attached to a trivalent black vertex separated by a single edge from a white vertex, define $s_1(G)$ to be

the graph obtained from G by moving λ to the white vertex. If G is not equivalent to a graph of this form, then define $s_1(G) = 0$. Then we have $d(s_1(G)) + s_1(d(G)) = G' - G$ where G' is obtained from G by moving λ_1 to the next half-edge attached at the white vertex. Indeed, the graphs G and G' are the “new” non-zero terms in $d(s_1(G))$. (In the example in Figure 7, the homotopy is a sum of 3 graphs, with p_7 , then p_2 , then p_4 attached in turn directly at the white vertex.) \square

Proof of Theorem 3.8. The proof of the theorem then follows directly using the same argument as for Theorem 3.1: the analogue of Lemma 3.5 holds, with Lemma 3.9 replacing Lemma 3.4, and this last result can then be used repeatedly to prove the theorem. \square

4. EXAMPLES OF NON-TRIVIAL HIGHER STRING TOPOLOGY OPERATIONS

We give in this section, as an example, two infinite families of cycles of increasing degree in the chain complex of Sullivan diagrams, associated to surfaces of increasing genus (resp. increasing number of boundary components), which induce a non-trivial action on the Hochschild homology of the cohomology algebra $H^*(S^n)$ for any $n \geq 2$, also over the rationals. As explained in the introduction, these operations can be used to define rational string topology operations for simply-connected manifolds using [22] and [16] (see also [9] or [36, Sec 6.6]), which are non-trivial as they act non-trivially on $HH_*(H^*(S^n), H^*(S^n)) \cong H^*(LS^n)$.

Once the classes are constructed, it will be easy to check that they represent cycles in the chain complex of Sullivan diagrams. The fact that they act non-trivially on the Hochschild homology of a particular algebra will then imply that they represent non-trivial homology classes. Very little is known about the homology of the chain complex of Sullivan diagrams. The method presented here can be used to detect additional families of classes, and gives us a first insight in the homology of this chain complex.

For each $g \geq 1$, we construct two black and white graphs μ_g and t_g as follows: The graph μ_g is obtained from the graph l_{2g+2} , a single white vertex with $2g + 2$ leaves, by adding a tree connecting the odd entries $1, 3, \dots, 2g + 1$ together and labeling the remaining leaves $1, 2, \dots, g + 1$ in their cyclic order. (See Figure 8 for a picture of μ_1, μ_2 and μ_3 both as the black and white graphs just described, and in the more classical picture of Sullivan chord diagrams using the isomorphism given in [36, Thm 5.9].)




FIGURE 8. The graphs t_g and μ_g for $g = 1, 2, 3$ as black and white graphs and as classical Sullivan diagrams

The graph t_g is then constructed from μ_g by adding a second tree connecting the remaining leaves at the white vertex, with a leaf on the new tree as shown in the figure for t_1, t_2 and t_3 .

The graphs μ_g and t_g both have degree $2g + 1$. It is easy to check that they have boundary 0: they have an even number of faces, and each face d_{2i} cancels with the face d_{2i+1} . Hence the graphs are cycle. The computation given below of a non-trivial action in homology using these graphs implies that they represent non-trivial homology classes.

We consider μ_g as living in $\mathcal{SD}([g+1]_0, [1]_0)$, that is each leaf is considered as marking an incoming closed boundary component. The topological type of the associated cobordism, which is the surface obtained by thickening the graph (with the white vertex defining a boundary component), is a surface of genus 0 with $g + 2$ boundary components, $g + 1$ of which being incoming and the last one being outgoing.

The graph t_g is considered as living in $\mathcal{SD}([1]_0, [1]_0)$. The surface obtained by thickening the has two boundary components, the incoming and the outgoing one, and an Euler characteristic computation shows that it has genus g . In the category of Sullivan diagrams, t_g is actually the precomposition of μ_g with a representative for the $g + 1$ -legged coproduct of degree 0 (the dual of the g -fold Chas-Sullivan product).

Consider the Frobenius algebra $A = \mathbb{Z}[x]/(x^2)$ concentrated in degree 0, with the coproduct given by $\nu(1) = 1 \otimes x + x \otimes 1$ and $\nu(x) = x \otimes x$. This is an $H_0(\mathcal{O})$ -algebra. (Up to degree shift and signs, this is the cohomology algebra of S^n .) The reduced Hochschild complex of A has two generating class in each degree, namely $1 \otimes x \otimes \cdots \otimes x$ and $x \otimes x \otimes \cdots \otimes x$, and the differential is 0 except on the terms $1 \otimes x \otimes \cdots \otimes x$ in degree $2i$, which have differential $2(x \otimes \cdots \otimes x)$ in degree $(2i - 1)$. In particular, the classes $1 \otimes x \otimes \cdots \otimes x$ in odd degree are cycles representing non-trivial homology classes in $HH_*(A, A)$, and so are the classes $x \otimes \cdots \otimes x$, though those living in odd degrees in this case only generate a $\mathbb{Z}/2$ in homology.

Proposition 4.1. *Let $A = \mathbb{Z}[x]/(x^2)$ be the $H_0(\mathcal{O})$ -algebra defined above and let*

$$F : \mathcal{OC}([1]_0, [1]_0) \otimes C_*(A, A) \longrightarrow C_*(A, A)$$

be the action of Sullivan diagrams of Theorem 3.8 on the Hochschild complex of A . Let $t_g \in \mathcal{SD}([1]_0, [1]_0)$, and $x \in C_0(A, A)$ be as above. Then

$$F(t_g \otimes x) = 1 \otimes x \otimes \cdots \otimes x \in C_{2g+1}(A, A).$$

In particular, $H_(F(t_g \otimes -)) : HH_0(A, A) \rightarrow HH_{2g+1}(A, A)$ is non-zero, also with rational coefficients.*

The following corollary follows from the fact that t_g can be expressed in Sullivan diagrams as the g -fold composition of the coproduct corresponding to the dual of the Chas-Sullivan product, which takes $x \in C_0(A, A)$ to $x \otimes \cdots \otimes x \in C_0(A, A)^{\otimes g+1}$, followed by μ_g , though the result can also be checked directly.

Corollary 4.2. *The class $\mu_g \in \mathcal{SD}([g+1]_0, [1]_0)$ induces a non-trivial map $HH_0(A, A)^{\otimes g+1} \rightarrow HH_{2g+1}(A, A)$ for $A = \mathbb{Z}[x]/(x^2)$ as in the proposition.*

Proof of Proposition 4.1. Indeed, following the recipe given in [36, 6.2] for reading of the action of a graph on an algebra, we put x at the leaf of t_g and read off the graph minus the white vertex as a composition of operations in A . In the case of t_g , this graph is the union of the two trees used to construct the graph above. The first tree, as an operation on A , is the iterated coproduct $\nu^g(1)$, and the second tree is an iterated coproduct $\nu^g(x) = x \otimes \cdots \otimes x$. We have $\nu^g(1) = 1 \otimes x \otimes \cdots \otimes x + x \otimes 1 \otimes x \otimes \cdots \otimes x + \cdots + x \otimes \cdots \otimes x \otimes 1$, but only the first term gives a non-zero result in the reduced Hochschild complex. The resulting Hochschild chain is an intertwine of the two iterated coproduct, as prescribed by the way the trees are attached to the white vertex. Explicitly, it is the class $1 \otimes x \otimes \cdots \otimes x$ with $2g + 1$ x 's, as announced.

For the statement about the action in homology, we just note that x and $1 \otimes x \otimes \cdots \otimes x$ both are cycles representing non-trivial homology class, each generating a free \mathbb{Z} -summand. \square

We consider briefly the case of the cohomology algebra $H^*(S^n)$, which is a degree shifted version of the algebra A above. Just as A it is generated by two classes: $1 \in H^0(S^n)$ and $x \in H^n(S^n)$, with 1 a unit for the product, and $x \cdot x = 0$. The coproduct ν has degree n and satisfies $\nu(1) = 1 \otimes x \pm x \otimes 1$ and $\nu(x) = x \otimes x$. (The sign does not

play a role here.) Again just as above, the reduced Hochschild complex of $H^*(S^n)$ is generated by a class $1 \otimes x \otimes \cdots \otimes x$, though now in total degree $in - i$ for each $i \geq 0$, and $x \otimes \cdots \otimes x$, now in total degree $in - i + 1$ for each $i \geq 1$. When n is odd, the differential of the complex is identically 0, and when n is even, it is 0 except in degrees $2in - 2i$, where it is multiplication by 2. In particular, the collection

$$1, 1 \otimes x, x, (1 \otimes x \otimes x), x \otimes x, 1 \otimes x \otimes x \otimes x, x \otimes x \otimes x, (1 \otimes x \otimes x \otimes x \otimes x), x \otimes x \otimes x \otimes x, \dots$$

ordered in increasing degree, represents a collection of generator of the Hochschild homology of $H^*(S^n)$, where the generators in bracket are only to be used for n is odd. Under the isomorphism $HH_{-*}(H^*(S^n), H^*(S^n)) \cong H^*(LM)$ for $n \geq 2$, the class $[x] \in HH_0(H^*(S^n), H^*(S^n))$ is dual to the n -dimensional class of constant loops coming from the inclusion $S^n \hookrightarrow LS^n$.

The graded vector space $H^*(S^n)$ is a dimension n Frobenius algebra in the sense of [36, Sec 6.4], and the action of Sullivan diagrams is replaced by a degree shifted version of the above action. The calculation goes through in the exact same way and there are no sign issues as signs did not play a role in the computation. Hence t_g and μ_g define non-trivial string topology operations.

5. APPENDIX A: THE CHAIN COMPLEX ASSOCIATED TO A COSIMPLICIAL SET

In this section, we show that the chain complex associated to a cosimplicial set has homology concentrated in degree 0, generated by the elements in cosimplicial degree 0 on which d_0 and d_1 agree. This result was suggested to us by Bill Dwyer. It can be found in alternative form in [28, Prop 23.10], [1, 2.2-2.4], [14, Lem 4.1].

We denote by Δ the category of finite sets $[n] = \{0, \dots, n\}$, $n \geq 0$, and monotone maps. A cosimplicial set is a functor $X^\bullet : \Delta \rightarrow Sets$. Recall that the coface map $d^i : X^q \rightarrow X^{q+1}$ is the image under X^\bullet of the injection, also denote $d^i : [q] \hookrightarrow [q+1]$, which omits i , and the codegeneracy $s^j : X^q \rightarrow X^{q-1}$ is the image of the surjection $s^j : [q] \twoheadrightarrow [q-1]$ mapping j and $j+1$ to j . Any injection in Δ is a composition of cofaces and any surjection is a composition of codegeneracies.

Lemma 5.1. *Let X^\bullet be a cosimplicial set, and $x \in X^q$ a q -simplex for some $q \geq 0$. There exists a unique $y \in \cup_{p \leq q} X^p$ so that $x = D(y)$ for $D = d^{i_1} \dots d^{i_{q-p}} \in \Delta(p, q)$ a composition of coface maps (i.e. an injection) and y not itself a coface. Moreover, the injection D is unique as a morphism in Δ unless $y \in X^0$ and $d^0 y = d^1 y$. In that last case, we have $Dy = D'y$ for any injection $D, D' \in \Delta(0, q)$.*

Proof. Suppose $x = D_1 y_1 = D_2 y_2$ for D_1, D_2 compositions of cofaces and $y_1 \in X^{p_1}$, $y_2 \in X^{p_2}$ which are not themselves cofaces. Let S_1 be a composition of codegeneracies (i.e. a surjection) such that $S_1 D_1 = id \in \Delta(p_1, p_1)$. Then $y_1 = S_1 D_1 y_1 = S_1 D_2 y_2$. Now write $S_1 D_2 = D_3 S_3 \in \Delta(p_2, p_1)$ for some D_3, S_3 compositions of cofaces and codegeneracies respectively—this is possible as any map in Δ can be written as a composition of a surjection followed by an injection. As y_1 is not a coface, we must have $D_3 = id$ and $y_1 = S_3 y_2$. By symmetry, we also have that $y_2 = S'_3 y_1$ for some composition of codegeneracies S'_3 . But this can only happen if $p_1 = p_2$, and $S_3 = S'_3 = id$. Hence $y_1 = y_2$.

For the uniqueness of D , suppose first that y is not in cosimplicial degree 0. As we had $S_1 D_1 = D_3 S_3$ in the above computation, with S_1 any left inverse of D_1 , we can conclude from the above that $S_1 D_2 = id$ for any left inverse S_1 of D_1 in the category Δ . But if $D_1 \neq D_2$, the corresponding maps in Δ are different, and if the source of D_1, D_2 is not 0, then there exists an S_1 such that $S_1 D_1 = id$ in Δ but $S_1 D_2 \neq id$.

We are left with the case y in degree 0. (Here we have that $S_1 D_1 = id$ for any $D_1 : [0] \hookrightarrow [q]$ and the unique surjection $S_1 : [q] \twoheadrightarrow [0]$.) Suppose first that $d^0 y = d^1 y$. We then prove by induction that $Dy = d^0 \dots d^0 y$ for any composition of coface maps. Indeed, suppose it is true for compositions of length $k-1$. Then write D of length k

as $D = d^i d^0 \dots d^0$. As $d^i d^0 = d^0 d^{i-1}$ for any $i > 0$, we have that $Dy = d^i d^0 \dots d^0 y = d^0 d^{i-1} d^0 \dots d^0 y = d^0 \dots d^0 y$.

Finally, if $d^0 y \neq d^1 y$, we can distinguish different compositions D_1, D_2 using the following fact: if $D_1, D_2 : [0] \rightarrow [q]$ are compositions of coface maps in Δ with $D_1(0) > D_2(0)$, there exists a composition of degeneracies $S : [q] \rightarrow [2]$ such that $SD_1 = d_0$ and $SD_2 = d_1$. Then we must have $D_1(y) \neq D_2(y)$ as for such a choice of S we have $SD_1(y) \neq SD_2(y)$. \square

By a *semi-cosimplicial set*, we mean a functor $Y^\bullet : \Delta^{inj} \rightarrow \mathit{Sets}$, where Δ^{inj} is the subcategory of Δ with the same objects but with only monotone injections as morphisms. So a semi-cosimplicial set has coface maps satisfying the usual relations, but no codegeneracies. Let $\mathit{Inj}(r)^\bullet$ be the semi-cosimplicial set represented by r , i.e. with $\mathit{Inj}(r)^q = \Delta^{inj}(r, q)$, and with the coface maps defined by composition. It is the free semi-cosimplicial set generated by one element in cosimplicial degree r .

The above lemma associates to every simplex x in a cosimplicial set X^\bullet , a unique minimal y such that $x = Dy$. Moreover, the lemma shows that, if $d^0 y \neq d^1 y$ then $Dy \neq D'y$ for any injection $D \neq D'$ in Δ , i.e. y generates a free semi-cosimplicial set in X^\bullet , and if $d_0 y = d_1 y$, then $Dy = D'y$ for all D, D' of the same length, i.e. y generates a constant sub-cosimplicial set (with y necessarily in degree 0). This proves the following:

Corollary 5.2. *Any cosimplicial set X^\bullet splits as a semi-cosimplicial set as the disjoint union of constant subcosimplicial sets, one for each $y \in X^0$ such that $d^0 y = d^1 y$, and free semi-cosimplicial sets $\mathit{Inj}(r)^\bullet$ for various r 's.*

To a (semi)-cosimplicial set X^\bullet , we can associate a chain complex $\mathbb{Z}X := \bigoplus_q \mathbb{Z}X^q$, with $\mathbb{Z}X^q$ in degree $-q$, and with differential $d_X = \sum (-1)^i d^i$, the alternating sum of the coface maps.

Proposition 5.3. *For any $r \geq 0$, the chain complex $(\mathbb{Z}\mathit{Inj}(r), d_{\mathit{Inj}})$ associated to the free semi-cosimplicial set $\mathit{Inj}(r)^\bullet$ has no homology.*

Proof. In degree q , $\mathbb{Z}\mathit{Inj}(r)$ is the free module on all injection $[r] \hookrightarrow [q]$, and the boundary map $d : \mathbb{Z}\mathit{Inj}(r)^q \rightarrow \mathbb{Z}\mathit{Inj}(r)^{q+1}$ is the alternating sum of the coface maps: $d = \sum_{i=1}^{q+1} (-1)^i d^i$, where $d^i([r] \xrightarrow{x} [q])$ is the composition $[r] \xrightarrow{x} [q] \xrightarrow{d_i} [q+1]$ where the second map does not hit the i th element.

Define a chain homotopy $s : \mathbb{Z}\mathit{Inj}(r)^q \rightarrow \mathbb{Z}\mathit{Inj}(r)^{q-1}$ by $s([r] \xrightarrow{x} [q])$ is $(-1)^q x$, considered as landing in $[q-1]$ if q is not in the image of x , and is 0 otherwise.

Then if q is in the image of x , we have $sd + ds([r] \xrightarrow{x} [q]) = s(\sum_{i=1}^{q+1} (-1)^i ([r] \xrightarrow{x} [q] \xrightarrow{d_i} [q+1])) + 0$ equals $(-1)^{q+1} (-1)^{q+1} (x)$ as only the last term in the sum will not hit $q+1$. On the other hand, if q is not in the image of x , then

$$\begin{aligned} sd + ds([r] \xrightarrow{x} [q]) &= s\left(\sum_{i=1}^{q+1} (-1)^i ([r] \xrightarrow{x} [q] \xrightarrow{d_i} [q+1])\right) \\ &\quad + (-1)^q \sum_{i=1}^q (-1)^i ([r] \xrightarrow{sx} [q-1] \xrightarrow{d_i} [q]) \end{aligned}$$

As x does not hit q , none of the terms in the first sum will hit $q+1$ and, given that s in this case introduces a sign $(-1)^{q+1}$, all but the last term will cancel with the terms in the second sum. The only term that does not cancel is equal to x . \square

We can now prove the result we use in Section 3:

Proposition 5.4. *Let X^\bullet be a cosimplicial set, and $C(X, M) := \bigoplus_q \mathbb{Z}X^q \otimes M$ the associated chain complex with trivial coefficients in a module M and differential $d_X =$*

$\sum (-1)^i d^i$. Then

$$H_*(C(X, M)) = \bigoplus_{\substack{x \in X^0 \\ d^0 x = d^1 x}} M$$

with each copy of M concentrated in degree 0.

Proof. By Corollary 5.2 the cosimplicial set X^\bullet splits as a semi-cosimplicial set into a disjoint union of free semi-cosimplicial sets $\text{Inj}(r)$ and one constant cosimplicial set for each $x \in X^0$ satisfying $d^0 x = d^1 x$. As the differential in $C(X, M)$ only uses the semi-cosimplicial set structure, we also get a splitting of $C(X, M)$ into such direct summands. By Proposition 5.3, the free semi-cosimplicial sets do not contribute anything to the homology. On the other hand, the chain complex associated to a constant cosimplicial set is the cochain complex of a point, and hence contributes a copy of the module M in degree 0. \square

6. APPENDIX B: GRAPH COMPLEXES

In this appendix, we briefly recall the necessary ingredients to define a chain complex of graphs as used in the paper. We exemplify how this allows to replace certain algebraic equations by graph equations.

A *graph* is formally defined as a tuple $G = (V, H, s, i)$ where V is its set of *vertices*, H its set of *half-edges*, $s : H \rightarrow V$ is the *source map*, and $i : H \rightarrow H$ is an involution. Unless otherwise stated, graphs will be assumed to be at least 3-valent, i.e. such that $|v| := |s^{-1}(v)| \geq 3$ for all $v \in V$. An *edge* in the graph is a pair of half-edges $(h, i(h))$, with $i(h) \neq h$. A *leaf* is a fixed point of the involution: $h = i(h)$. We allow also the exceptional graph made out of a single leaf and no vertices, for example to describe identity morphisms in e.g. the category \mathcal{A}_∞ . An *orientation* for a graph G is a unit vector in $\det(\mathbb{R}(V \sqcup H))$.

To a graph G , one can associate a 1-dimensional CW-complex, its *realization*, by attaching edges and leaves along their end vertices, in the way prescribed by the source map. A *tree* is a graph such that its realization is contractible. A *planar tree* is a tree, together with the additional data of an embedding of the realization of the tree into the plane.

A *fat graph* is a graph equipped with a cyclic ordering of the half-edges $s^{-1}(v)$ at each vertex v . Planar graphs are examples of fat graphs, the embedding in the plane giving such a cyclic ordering at each vertex.

Given a graph $G = (V, H, s, i)$ and an edge $e = (h_1, h_2)$ of G with $s(h_1) \neq s(h_2)$, one can define the collapse G/e of G along e by removing the two half-edges h_1, h_2 and identifying their source vertices. If G had a fat structure, G/e inherits a fat structure. If G is oriented, G/e inherits an orientation as follows: writing the orientation of G as $v_1 \wedge v_2 \wedge h_1 \wedge h_2 \wedge X$ for $v_1 = s(h_1)$ and $v_2 = s(h_2)$, we defined the orientation of G/e to be $v \wedge X$ where v denotes the vertex obtained by identifying v_1 and v_2 . Now a *blow-up* of an (oriented, fat) graph G is an (oriented, fat) graph \hat{G} such that there exists an edge e in \hat{G} with $\hat{G}/e \cong G$ as (oriented, fat) graphs.

Blow-ups can be used to define a chain complex of oriented fat graphs as follows. First we define the *degree* of a graph G as $\text{deg}(G) = \sum_{v \in V} |v| - 3$. Now one defines a chain complex which, in degree d , is the free \mathbb{Z} -module on the set of oriented degree d fat graphs, modulo the relation that (-1) reverses the orientation. The *differential* is defined as the sum of all possible blow-ups:

$$d(G) = \sum_{\substack{(\hat{G}, e) \\ \hat{G}/e \cong G}} \hat{G}$$

(See Figure 9 for an example.)

Graph complexes are used in the paper to describe \mathcal{A}_∞ -algebras, compute their Hochschild complex and compute operations on the Hochschild complex. We give here an example on how to go between graphs and algebra, and how graph orientations translate to signs. Hopefully it illustrates that, in an appropriate context, oriented graphs can be easier to work with than algebraic formulas with signs.

Example 6.1 (The map $m_2 \in \mathcal{A}_\infty(2, 1)$ is homotopy associative.). Let m_2 be the graph with one vertex v and three leaves h_0, h_1, h_2 attached to it. (So $s(h_i) = v$ and $i(h_i) = h_i$ for all i .) We give m_2 the “left-right top-down” orientation $h_1 \wedge h_2 \wedge v \wedge h_0$, and we interpret m_2 as a morphism from 2 to 1 in \mathcal{A}_∞ by labeling h_1 and h_2 as the first and second incoming leaves, and h_0 as the root/outgoing leaf. Similarly, let m_3 be the graph with a single vertex w and leaves l_0, l_1, l_2, l_3 attached to it, with orientation $l_1 \wedge l_2 \wedge l_3 \wedge w \wedge l_0$.

The graph m_3 has degree 1. Its differential has two terms, shown in Figure 9, as there are two ways of blowing w up. Both blow-ups can be described as having vertex set $V = \{w_1, w_2\}$, half-edges $\{l_0, l_1, l_2, l_3, e_1, e_2\}$ with $i(e_1) = e_2$, $s(e_i) = w_i$, though with different source maps for the half-edges l_i , as described by the picture. To compute their orientation, we first write $l_1 \wedge l_2 \wedge l_3 \wedge w \wedge l_0 = -w \wedge l_1 \wedge l_2 \wedge l_3 \wedge l_0$. Now by definition of the differential, both terms come with the orientation $-w_1 \wedge w_2 \wedge e_1 \wedge e_2 \wedge l_1 \wedge l_2 \wedge l_3 \wedge l_0$.

The oriented graph m_3 defines an element in $\mathcal{A}_\infty(3, 1)$ with l_1, l_2, l_3 the three incoming leaves and l_0 the outgoing leaf. We would like to compare the terms in its differential to the compositions $m_2 \circ (m_2 + id)$ and $m_2 \circ (id + m_2)$. As graphs, these are indeed two terms, but the orientation of $m_2 \circ (m_2 + id)$ is $(l_1 \wedge l_2 \wedge w_1 \wedge e_1) \wedge (e_2 \wedge l_3 \wedge w_2 \wedge l_0)$ and that of $m_2 \circ (id + m_2)$ is $(l_2 \wedge l_3 \wedge w_1 \wedge e_1) \wedge (l_1 \wedge e_2 \wedge w_2 \wedge l_0)$, the orientation of a composition being the juxtaposition of the orientations. One then checks that, with the above choices of oriented representatives for m_2, m_3 , we have $d(m_3) = m_2 \circ (m_2 + id) - m_2 \circ (id + m_2)$.




FIGURE 9. The differential of m_3

More generally, the left-right top-bottom orientation gives a “canonical” orientation for planar rooted trees (or trees with a leaf specified as output) and this can be used to translate any formula given in terms of oriented graphs in \mathcal{A}_∞ as an algebraic formula in terms of m_i ’s. Likewise, one can choose a canonical orientation for the graph l_k of Section 1 and translate graph equations for the Hochschild complex of a dg-algebra to algebraic equations.

We refer the reader to [36], in particular Sections 2,3 and 7 for more details and further examples of graph complexes modeling algebraic structures.

REFERENCES

- [1] A. K. Bousfield. On the homology spectral sequence of a cosimplicial space. *Amer. J. Math.*, 109(2):361–394, 1987.
- [2] Henri Cartan and Samuel Eilenberg. *Homological algebra*. Princeton University Press, Princeton, N. J., 1956.
- [3] Ralph L. Cohen and Véronique Godin. A polarized view of string topology. In *Topology, geometry and quantum field theory*, volume 308 of *London Math. Soc. Lecture Note Ser.*, pages 127–154. Cambridge Univ. Press, Cambridge, 2004.

- [4] K. Costello. Topological conformal field theories and Calabi-Yau categories. *Adv. Math.*, 210(1):165–214, 2007.
- [5] Kevin Costello. A dual version of the ribbon graph decomposition of moduli space. *Geom. Topol.*, 11:1637–1652, 2007.
- [6] Yukio Doi. Homological coalgebra. *J. Math. Soc. Japan*, 33(1):31–50, 1981.
- [7] Daniela Egas. Comparing fat graph models of moduli space. Preprint in preparation.
- [8] Daniela Egas and Alexander Kupers. Comparing fat graph models of moduli space and their compactifications. Preprint (2014).
- [9] Yves Félix and Jean-Claude Thomas. Rational BV-algebra in string topology. *Bull. Soc. Math. France*, 136(2):311–327, 2008.
- [10] Benoit Fresse. Koszul duality of operads and homology of partition posets. In *Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory*, volume 346 of *Contemp. Math.*, pages 115–215. Amer. Math. Soc., Providence, RI, 2004.
- [11] Benoit Fresse. *Modules over operads and functors*, volume 1967 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2009.
- [12] Véronique Godin. Higher string topology operations. Preprint 2007. (arXiv:0711.4859).
- [13] Véronique Godin. The unstable integral homology of the mapping class groups of a surface with boundary. *Math. Ann.*, 337(1):15–60, 2007.
- [14] Thomas G. Goodwillie. A remark on the homology of cosimplicial spaces. *J. Pure Appl. Algebra*, 127(2):167–175, 1998.
- [15] J. F. Jardine. Cosimplicial spaces and cocycles. Preprint 2011.
- [16] John D. S. Jones. Cyclic homology and equivariant homology. *Invent. Math.*, 87(2):403–423, 1987.
- [17] Angela Klamt. The complex of formal operations on the Hochschild chains of commutative algebras. Preprint 2013 (arXiv:1309.7882).
- [18] Angela Klamt. The complex of looped diagrams and natural operations on Hochschild homology. PhD thesis, 2013 (available at <http://www.math.ku.dk/noter/filer/phd13ak.pdf>).
- [19] Angela Klamt. Natural operations on the Hochschild complex of commutative Frobenius algebras via the complex of looped diagrams. Preprint 2013 (arXiv:1309.4997).
- [20] Joachim Kock. *Frobenius algebras and 2D topological quantum field theories*, volume 59 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2004.
- [21] M. Kontsevich and Y. Soibelman. Notes on A_∞ -algebras, A_∞ -categories and non-commutative geometry. In *Homological mirror symmetry*, volume 757 of *Lecture Notes in Phys.*, pages 153–219. Springer, Berlin, 2009.
- [22] Pascal Lambrechts and Don Stanley. Poincaré duality and commutative differential graded algebras. *Ann. Sci. Éc. Norm. Supér. (4)*, 41(4):495–509, 2008.
- [23] Aaron D. Lauda and Hendryk Pfeiffer. Open-closed strings: two-dimensional extended TQFTs and Frobenius algebras. *Topology Appl.*, 155(7):623–666, 2008.
- [24] Jean-Louis Loday. *Cyclic homology, second ed.*, volume 301 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1998.
- [25] Jean-Louis Loday and Bruno Valette. *Algebraic Operads*, volume 346 of *Grundlehren der mathematischen Wissenschaften*.
- [26] Saunders Mac Lane. Categorical algebra. *Bull. Amer. Math. Soc.*, 71:40–106, 1965.
- [27] I. Macdonald. *Symmetric functions and Hall polynomials (second edition)*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 1995.
- [28] J. Peter May. *Simplicial objects in algebraic topology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1992. Reprint of the 1967 original.
- [29] Ryszard Nest and Boris Tsygan. On the cohomology ring of an algebra. In *Advances in geometry*, volume 172 of *Progr. Math.*, pages 337–370. Birkhäuser Boston, Boston, MA, 1999.
- [30] R. C. Penner. Lambda lengths. preprint (2006), available at http://www.ctqm.au.dk/research/MCS/lambda_lengths.pdf.
- [31] R. C. Penner. Decorated Teichmüller theory of bordered surfaces. *Comm. Anal. Geom.*, 12(4):793–820, 2004.
- [32] Teimuraz Pirashvili. Hodge decomposition for higher order Hochschild homology. *Ann. Sci. École Norm. Sup. (4)*, 33(2):151–179, 2000.
- [33] Jim Stasheff. Homotopy associativity of h-spaces i & ii. *Trans. Amer. Math. Soc.*, 108:275–292, 293–312, 1963.
- [34] Hirotaka Tamanoi. Stable string operations are trivial. *Int. Math. Res. Not. IMRN*, (24):4642–4685, 2009.
- [35] Thomas Tradler and Mahmoud Zeinalian. On the cyclic Deligne conjecture. *J. Pure Appl. Algebra*, 204(2):280–299, 2006.
- [36] N. Wahl and C. Westerland. Hochschild homology of structured algebras. Preprint 2011. (arXiv:1110.0651).

- [37] Charles A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.

E-mail address: `wahl@math.ku.dk`