

FROM MAPPING CLASS GROUPS TO AUTOMORPHISM GROUPS OF FREE GROUPS

NATHALIE WAHL

ABSTRACT

It is shown that the natural map from the mapping class groups of surfaces to the automorphism groups of free groups induces an infinite loop map on the classifying spaces of the stable groups after plus construction. The proof uses automorphisms of free groups with boundaries which play the role of mapping class groups of surfaces with several boundary components.

1. Introduction

Both the stable mapping class group of surfaces Γ_∞ and the stable automorphism group of free groups Aut_∞ give rise to infinite loop spaces $\text{B}\Gamma_\infty^+$ and BAut_∞^+ when taking the plus-construction of their classifying spaces. By the work of Madsen and Weiss [16], $\text{B}\Gamma_\infty^+$ is now well understood, whereas BAut_∞^+ remains rather mysterious. In this paper, we relate these two spaces by showing that the natural map $\text{B}\Gamma_\infty^+ \rightarrow \text{BAut}_\infty^+$ is a map of infinite loop spaces. This means in particular that the map $H_*(\Gamma_\infty) \rightarrow H_*(\text{Aut}_\infty)$ respects the Dyer–Lashof algebra structure. To prove this result, we introduce a new family of groups, the automorphism groups of free groups with boundary, which have the same stable homology as the automorphisms of free groups but enable us to define new operations.

Let F_n be the free group on n generators, and let $\text{Aut}(F_n)$ be its automorphism group. Note that $\text{Aut}(F_n) \cong \pi_0 \text{Htpy}_*(\vee_n S^1)$, the group of components of the pointed self-homotopy equivalences of a wedge of n circles. A result of Hatcher and Vogtmann [11] says that the inclusion map $\text{Aut}(F_n) \rightarrow \text{Aut}(F_{n+1})$ induces an isomorphism $H_i(\text{Aut}(F_n)) \rightarrow H_i(\text{Aut}(F_{n+1}))$ when $i \leq (n-3)/2$, and hence the homology of the stable automorphism group $\text{Aut}_\infty := \text{colim}_{n \rightarrow \infty} \text{Aut}(F_n)$ carries information about the homology of $\text{Aut}(F_n)$ for n large enough.

Let $S_{g,k}$ be a surface of genus g with k boundary components, and let $\Gamma_{g,k} := \pi_0 \text{Diff}^+(S_{g,k}; \partial)$ be its mapping class group, the group of components of the orientation-preserving diffeomorphisms which fix the boundary pointwise. The maps $\Gamma_{g,k} \rightarrow \Gamma_{g+1,k}$, induced by gluing a torus with two discs removed, and $\Gamma_{g,k} \rightarrow \Gamma_{g,k+1}$, induced by gluing a pair of pants, are homology isomorphisms in dimension $i \leq (g-1)/2$ by Harer and Ivanov [9, 14]. Let $\Gamma_\infty := \text{colim}_{g \rightarrow \infty} \Gamma_{g,1}$ denote the stable mapping class group.

There is a map $f: \Gamma_{g,1} \rightarrow \text{Aut}(F_{2g})$ obtained by considering the action on the fundamental group of the surface $S_{g,1}$ since $\pi_1(S_{g,1}) \cong F_{2g}$. The spaces $\coprod_{g \geq 0} \text{B}\Gamma_{g,1}$ and $\coprod_{n \geq 0} \text{BAut}(F_n)$ have monoid structures induced by the pair of pants

Received 8 July 2004; revised 16 December 2004.

2000 *Mathematics Subject Classification* 55P47 (primary), 19D23, 20F28 (secondary).

This work was supported by a Marie Curie Fellowship of the European Community under contract number HPMF-CT-2002-01925.

multiplication on surfaces $S_{g,1} \times S_{h,1} \rightarrow S_{g+h,1}$ and by wedging circles $\vee_n S^1 \times \vee_m S^1 \rightarrow \vee_{n+m} S^1$ respectively. The map $Bf: \coprod_{g \geq 0} B\Gamma_{g,1} \rightarrow \coprod_{n \geq 0} \text{BAut}(F_n)$ is a map of monoids, and hence we have a map of loop spaces $\mathbb{Z} \times B\Gamma_\infty^+ \rightarrow \mathbb{Z} \times \text{BAut}_\infty^+$ on the group completion. The wedge product defines an infinite loop structure on $\mathbb{Z} \times \text{BAut}_\infty^+$ as it defines a symmetric monoidal structure on $\coprod \text{Aut}(F_n)$, thought of as a category. The infinite loop structure on $\mathbb{Z} \times B\Gamma_\infty^+$, discovered by Tillmann [22], is more complicated; Tillmann defines a cobordism 2-category \mathcal{S} which is symmetric monoidal under disjoint union and such that $\Omega B\mathcal{S} \simeq \mathbb{Z} \times B\Gamma_\infty^+$. As $B\mathcal{S}$ is an infinite loop space, so is its loop space $\mathbb{Z} \times B\Gamma_\infty^+$. Our main result is the following.

THEOREM 1.1. *There is an infinite loop space structure on $\mathbb{Z} \times \text{BAut}_\infty^+$ equivalent to the one induced by wedging circles and such that the map*

$$\mathbb{Z} \times B\Gamma_\infty^+ \rightarrow \mathbb{Z} \times \text{BAut}_\infty^+$$

induced by the action on the fundamental group is a map of infinite loop spaces.

To prove this theorem, we enlarge Tillmann’s cobordism category \mathcal{S} by introducing an extra ‘graph-like’ morphism. We obtain a new 2-category \mathfrak{T} which contains \mathcal{S} as a subcategory and such that $\Omega B\mathfrak{T} \simeq \mathbb{Z} \times \text{BAut}_\infty^+$. The theorem then follows from the fact that the inclusion $\mathcal{S} \rightarrow \mathfrak{T}$ is a map of symmetric monoidal categories.

The category \mathfrak{T} is closely related to the *automorphism groups of free groups with boundary*, which we define now.

Let $G_{n,k}$ be the graph shown in Figure 1 consisting of a wedge of n circles together with k extra circles joined by edges to the basepoint. We call the k circles disjoint from the basepoint *boundary circles*. The automorphism group of free group with boundary $A_{n,k}$ is by definition $\pi_0 \text{Htpy}(G_{n,k}; \partial)$, the group of components of the space of homotopy equivalences of $G_{n,k}$ which fix the basepoint and the k boundary circles pointwise. In particular, $A_{n,0} = \text{Aut}(F_n)$. The group $A_{n,k}$ is an analogue of $\Gamma_{g,k+1}$. In fact, $\Gamma_{g,k+1}$ is a subgroup of $A_{2g,k}$. A description of $A_{n,k}$ as an extension of a subgroup of $\text{Aut}(F_{n+k})$, as well as a presentation of the group, are given in joint work with Jensen [15] and a description in terms of mapping class groups of certain 3-manifolds is given in joint work with Hatcher [13]. This last description is used in [13] to prove that the natural inclusions $A_{n,k} \rightarrow A_{n+1,k}$ and $A_{n,k} \rightarrow A_{n,k+1}$ are homology isomorphisms in dimension $i \leq (n - 3)/3$.




FIGURE 1. $G_{n,k}$.

The boundary circles of $G_{n,k}$ allow to define new gluing operations between graphs and between graphs and surfaces. We use these operations to define the 2-category \mathfrak{T} , whose objects are the natural numbers, whose 1-morphisms are built out of graphs and surfaces by gluing and disjoint union, and whose 2-morphisms are homotopy equivalences fixing the boundary. The cobordism 2-category \mathcal{S} of [22] is the subcategory of \mathfrak{T} generated by the 1-morphisms built out of surfaces only. (See Section 2.)

Theorem 4.1 says that $\Omega B\mathfrak{T} \simeq \mathbb{Z} \times \mathrm{BAut}_\infty^+$. The main ingredients of the proof are the homological stability of the automorphisms of free groups with boundary and a generalized group completion theorem. Our Theorem 5.1 then says that the infinite loop space structure on $\mathbb{Z} \times \mathrm{BAut}_\infty^+$ induced by the symmetric monoidal structure of \mathfrak{T} is equivalent to the one previously known, induced by wedging circles.

Tillmann proved in [23] that the map $\mathbb{Z} \times \mathrm{B}\Gamma_\infty^+ \rightarrow A(*)$ to Waldhausen's space $A(*)$ is a map of infinite loop spaces. Our result says that this map factors as a map of infinite loop spaces: $\mathbb{Z} \times \mathrm{B}\Gamma_\infty^+ \rightarrow \mathbb{Z} \times \mathrm{BAut}_\infty^+ \rightarrow A(*)$. The space QS^0 factors out of each of these three spaces (away from two in the case of the mapping class group) and the first map on the QS^0 factor is multiplication by 2. The work of Dwyer–Weiss–Williams ([4], see also [23]) implies that the composite map $\mathbb{Z} \times \mathrm{B}\Gamma_\infty^+ \rightarrow A(*)$ factors through QS^0 . It is unknown whether $\mathbb{Z} \times \mathrm{B}\Gamma_\infty^+ \rightarrow \mathbb{Z} \times \mathrm{BAut}_\infty^+$ already factors through QS^0 , or for that matter whether $\mathbb{Z} \times \mathrm{BAut}_\infty^+$ is QS^0 . The rational homology of Aut_∞ is known to be trivial up to dimension 6 and is conjecturally trivial in all dimensions [12]. On the other hand, the homology of the mapping class group is known with any field coefficients [7] and is rather rich.

The paper is organized as follows. We construct the 2-category \mathfrak{T} in Section 2. In Section 3, we define a \mathfrak{T} -diagram which is used in Section 4 to show that $\Omega B\mathfrak{T} \simeq \mathbb{Z} \times \mathrm{BAut}_\infty^+$. In Section 5, we show the equivalence of the two infinite loop space structures on $\mathbb{Z} \times \mathrm{BAut}_\infty^+$. Section 6 gives an alternative definition of \mathfrak{T} using punctured surfaces.

2. Cobordism category with graphs

The $(1 + 1)$ -cobordism 2-category has objects 1-dimensional manifolds, 1-morphisms cobordisms between these manifolds and 2-morphisms diffeomorphisms of cobordisms restricting to the identity on the boundary. We define here a 2-category \mathfrak{T} , modifying Tillmann's model of the cobordism category in by adding an extra 1-morphism from the circle to itself, and replacing diffeomorphisms by homotopy equivalences.

2.1. Objects

The objects of \mathfrak{T} are the natural numbers $n \in \mathbb{N}$, where n can be thought of as a disjoint union of n circles.

2.2. 1-morphisms

To define the 1-morphisms of \mathfrak{T} , consider the following directed building blocks, called *pieces*:

- (1) a pair of pants \mathbb{P} , a torus with two discs removed \mathbb{T} and a disc \mathbb{D} with respectively 2, 1 and 0 incoming boundary components and each with 1 outgoing boundary;




FIGURE 2. Pieces and example of a 1-morphism from 4 to 2.

- (2) a graph-like piece $\mathbb{G} \simeq G_{1,1} \vee D^2$, with incoming boundary the boundary circle of $G_{1,1}$ and outgoing boundary the boundary of the disc (see Figure 2).

We fix a parametrization $[0, 2\pi[$ of the boundary circles, with 0 at the end point of the attaching edge for the incoming boundary of \mathbb{G} . We then allow the following gluing operation on the pieces: an incoming circle of one piece can be identified (using the parametrization) with the outgoing circle of another piece. The boundary of the glued object is defined to be the union of the boundary of the two pieces, minus the two identified circles.

A 1-morphism in \mathfrak{T} from n to m is a couple (T, σ) , where T is a 2-dimensional CW-complex obtained from the above pieces by gluing and disjoint union, with n incoming and m outgoing boundary components, and σ is a labelling of these boundaries. (See Figure 2 for an example.) One can moreover take disjoint union with copies of the circle, thought of as a morphism from 1 to 1. These circles are thus to be labelled on both sides. The $n!$ morphisms from n to n given by n disjoint copies of the circle correspond to the permutations of the labels, with the identity permutation representing the identity morphism on n .

One should think of (T, σ) as a combinatorial object, to which a topological space is associated.

For simplicity, we will drop σ from the notation in (T, σ) . Note that the building blocks of \mathfrak{T} are defined in such way that a 1-morphism $T : n \rightarrow m$ has exactly m connected components.

2.3. 2-morphisms

A 2-category is a category enriched over categories and the 1-morphisms of \mathfrak{T} are the objects of the categories of morphisms $\mathfrak{T}(n, m)$. The 2-morphisms are the morphisms in these categories. Given two 1-morphisms T and T' , objects of $\mathfrak{T}(n, m)$, the set of 2-morphisms between T and T' is $\pi_0 \text{Htpy}(T, T'; \partial)$, the group of components of the space of homotopy equivalences from T to T' which fix the boundary, that is which map the boundary of T to the boundary of T' via the identity map according to the labels. By [10, Proposition 0.19], any homotopy equivalence $T \rightarrow T'$ which fixes the boundary is a homotopy equivalence relative to the boundary. (The pair $(T, \partial T)$ is a CW-pair and hence satisfies the homotopy extension property, in which case the proposition applies.) In particular, every 2-morphism is invertible, but more importantly we will be able to glue homotopy equivalences along the boundaries.

REMARK 2.1. We consider homotopy classes of homotopy equivalences relative to the boundary. By homotopy classes, we mean path components in $\text{Htpy}(T, T'; \partial)$. Two elements f and g are in the same path component if and only if they are homotopy equivalent *relative to the boundary*. Note that here, it would not be equivalent to forget the ‘relative to the boundary’!

Note that if T and T' are permutations, the set of 2-morphisms is empty unless $T = T'$ in which case it is just the identity. If T and T' are surfaces, that is if they are built out of \mathbb{P} , \mathbb{T} and \mathbb{D} only, then

$$\pi_0 \text{Htpy}(T, T'; \partial) \cong \pi_0 \text{Diff}^+(T, T'; \partial)$$

(see [15, Section 2]). We describe in Lemma 2.2 below the 1-morphisms with automorphism group given by automorphisms of free groups with boundaries.

2.4. *Composition*

The composition of 1-morphisms is defined by gluing according to the labels. We will denote this composition by $T_1 \square T_2 : n \rightarrow p$ for $T_1 : n \rightarrow m$ and $T_2 : m \rightarrow p$.

For $f_1 : T_1 \rightarrow T'_1$ and $f_2 : T_2 \rightarrow T'_2$, define the horizontal composition $f_1 \square f_2 : T_1 \square T_2 \rightarrow T'_1 \square T'_2$ by applying f_1 to T_1 and f_2 to T_2 . This defines a homotopy equivalence as f_1 and f_2 are homotopy equivalences relative to the boundary.

Finally, vertical composition of 2-morphisms is given by the composition of homotopy equivalences.

All compositions are associative and this defines a 2-category. We write all compositions in \mathfrak{X} in the left-to-right order.

We have seen that surface 1-morphisms in \mathfrak{X} have automorphism group given by mapping class groups. We now describe the 1-morphisms with automorphism group given by automorphisms of free groups with boundary. They play an important role in determining the homotopy type of \mathfrak{X} .

Let $G_{n,k}$ be the graph given in Figure 1 and consider $G_{n,k} \vee D^2$, where the basepoint of the disc is its centre. The boundary of $G_{n,k} \vee D^2$ consists of $k + 1$ circles: the k boundary circles of $G_{n,k}$ together with the boundary circle of the disc D^2 . The graph $G_{0,0}$ is just a point so $G_{0,0} \vee D^2 \cong \mathbb{D}$. Note also that $G_{1,1} \vee D^2 = \mathbb{G}$.

Recall that $A_{n,k} = \pi_0 \text{Htpy}(G_{n,k}; \partial)$, where $\partial G_{n,k}$ is the basepoint of $G_{n,k}$ union its k boundary circles.

We call a *surface component* of an object T of $\mathfrak{X}(k, 1)$ any connected component of surface pieces $\mathbb{P}, \mathbb{T}, \mathbb{D}$ and the disc of \mathbb{G} glued together in T . For example, the object of $\mathfrak{X}(2, 1)$ in Figure 3 has two surface components: a torus with three holes and a disc. The *outgoing surface component* of T is the surface component whose outgoing boundary is the outgoing boundary of T . In the example, it is a disc.




FIGURE 3. $G_{3,2} \vee D^2$ homotopic to an object of $\mathfrak{X}_D(2, 1)$.

LEMMA 2.2. *An object T of $\mathfrak{T}(k, 1)$ is homotopic relative to the boundary to $G_{n,k} \vee D^2$ for some n , if and only if the outgoing surface component of T is a disc. Moreover, if this is the case, we have $\pi_0 \text{Htpy}(T; \partial) \cong A_{n,k}$.*

Let $\mathfrak{T}_D(k, 1)$ be the full subcategory of $\mathfrak{T}(k, 1)$ generated by the objects with a disc as outgoing surface component, and let $\mathfrak{T}_S(k, 1)$ be the full subcategory of $\mathfrak{T}(k, 1)$ generated by the other objects, that is the objects with outgoing surface component of higher genus or with more boundary components, or the object represented by the circle in $\mathfrak{T}(1, 1)$. By the lemma, there can be no morphisms in $\mathfrak{T}(k, 1)$ between the objects of $\mathfrak{T}_D(k, 1)$ and the objects of $\mathfrak{T}_S(k, 1)$. Hence we have the following.

COROLLARY 2.3.

$$\mathfrak{T}(k, 1) = \mathfrak{T}_D(k, 1) \coprod \mathfrak{T}_S(k, 1) \quad \text{and} \quad \text{B}\mathfrak{T}_D(k, 1) \simeq \coprod_{n \geq \varepsilon} \text{BA}_{n,k},$$

where $\varepsilon = 0$ when $k = 0$ and $\varepsilon = 1$ otherwise.

Proof of Lemma 2.2. Every surface component of an object T has exactly one outgoing boundary by construction. Suppose that T is an object of $\mathfrak{T}(k, 1)$ with outgoing surface component a disc. Then the outgoing boundary of any other surface component in T will be free, that is not fixed by the homotopy equivalences – as it cannot be the outgoing boundary of T . One can thus homotope each of these surface components S to a graph $G_{2m,l}$ lying in S , with basepoint 0 of the outgoing boundary of S , its l boundary circles on the l incoming boundaries of S and where m is the genus of S , and this can be done while keeping the boundary of T fixed. Such a homotopy can easily be constructed by considering the surface component as a polygon with appropriate identifications of the edges and with $l + 1$ discs removed.

Suppose on the other hand that T does not end with a disc component and let S be the outgoing surface component of T . If S has incoming boundary components, then they have to be incoming boundaries of T because the only non-surface piece one can glue to S is \mathbb{G} and this closes the incoming boundary it is glued to. Hence all the boundary components of S are boundary components of T and S cannot be homotoped to a graph while fixing the boundary of T . (Recall that $\pi_0 \text{Htpy}(S; \partial) \cong \pi_0 \text{Diff}^+(S; \partial)$ for a surface S .)

Finally, we want to see that if T is homotopic to $G_{n,k} \vee D^2$ relative to the boundary, then $\pi_0 \text{Htpy}(T; \partial) \cong A_{n,k}$. It is equivalent to show that $\pi_0 \text{Htpy}(G_{n,k} \vee D^2; \partial) \cong A_{n,k}$. Let $X := G_{n,k} \vee D^2$.

Note first that $A_{n,k} \cong \pi_0 \text{Htpy}(X; \tilde{\partial})$, where $\tilde{\partial}X$ consists of the k boundary circles of $G_{n,k}$ together with the basepoint $* = 0 \in \partial D^2$. There is an inclusion

$$\text{Htpy}(X; \partial) \hookrightarrow \text{Htpy}(X; \tilde{\partial})$$

and we want to show that it induces an isomorphism on π_0 .

Surjectivity: Let $f \in \text{Htpy}(X; \tilde{\partial})$. Then f maps ∂D^2 to a trivial loop in X . It is then easy to homotope f relative to $\tilde{\partial}X$ to a map fixing ∂D^2 . (One can also use the homotopy extension property of $(X, \partial X)$.)

Injectivity: Suppose that $f, g \in \text{Htpy}(X; \partial)$ are homotopic relative to $\tilde{\partial}$ by a homotopy $H : X \times I \rightarrow X$. Consider the restriction of H

$$H_{\partial D^2} : \partial D^2 \times I \rightarrow X.$$

Let $\hat{H}_{\partial D^2}$ be a lift of $H_{\partial D^2}$ to the universal cover of X , which is contractible. As $H_{\partial D^2}(0 \times I) = *$ and $H_{\partial D^2}(\partial D^2 \times 0) = H_{\partial D^2}(\partial D^2 \times 1) = \partial D^2$, we have $\hat{H}_{\partial D^2}(\partial D^2 \times 0) = \hat{H}_{\partial D^2}(\partial D^2 \times 1)$ is a fixed lift $\partial \hat{D}^2$ of ∂D^2 . Hence $\hat{H}_{\partial D^2}$ is homotopic to the constant map $i_{\partial D^2} : \partial D^2 \times I \rightarrow \partial \hat{D}^2$ relative to $\partial D^2 \times \{0, 1\}$. Let $h : (\partial D^2 \times I) \times I \rightarrow X$ be the projection of this last homotopy and extend it to $h_\partial : ((\partial X \times I) \cup (X \times \{0, 1\})) \times I \rightarrow X$ using the constant map on $\partial X \setminus \partial D^2$ and f and g on $(X \times 0) \times I$ and $(X \times 1) \times I$. Hence h_∂ is a homotopy from the restriction H_∂ of H to $(\partial X \times I) \cup (X \times \{0, 1\})$ to the map $i_{\partial X} \cup f \cup g : (\partial X \times I) \cup (X \times \{0, 1\}) \rightarrow X$. Now $(X \times I, (\partial X \times I) \cup (X \times \{0, 1\}))$ has the homotopy extension property as it is a CW-pair, so we get a homotopy $\mathbf{h} : (X \times I) \times I \rightarrow X$ from H to a homotopy $H' : X \times I \rightarrow X$ with $H'(X \times 0) = f(X)$, $H'(X \times 1) = g(X)$ and $H'(\partial X \times I)$ is the constant map on ∂X . Hence f and g are homotopic relative to ∂X . \square

By a Δ -category, we mean a category enriched over simplicial sets.

Let \mathcal{T} denote the Δ -category obtained from \mathfrak{T} by taking the nerve of the categories of morphisms, that is \mathcal{T} has the same objects as \mathfrak{T} , and $\mathcal{T}(n, m) := N_\bullet \mathfrak{T}(n, m)$.

PROPOSITION 2.4. *The Δ -category \mathcal{T} is symmetric monoidal under disjoint union and its classifying space $B\mathcal{T}$ is an infinite loop space.*

Proof. Taking disjoint union $T_1 \sqcup T_2$ and shifting the labels of T_2 induces a monoidal structure on \mathcal{T} . The symmetries are then given by the block permutations $n + m \rightarrow m + n$ in $\mathcal{T}(n + m, n + m)$.

Now $B\mathcal{T}$ is connected as there is a morphism from any object to 1. It follows that $B\mathcal{T}$ is an infinite loop space [17, 21]. \square

The cobordism Δ -category \mathcal{S} defined in [23] is the subcategory of \mathcal{T} generated by the surface pieces \mathbb{P}, \mathbb{T} and \mathbb{D} , that is it has the same objects as \mathcal{T} and the morphism space $\mathcal{S}(n, m)$ is the nerve of the full subcategory of $\mathfrak{T}(n, m)$ generated by the surface objects. (To be precise, the above model for \mathcal{S} is a mix of [22] and [23]: we use the combinatorial description of [23] but work with mapping class groups as in [22], instead of spaces of diffeomorphisms. This is equivalent because the components of the diffeomorphism group of a surface are contractible when the genus is large enough [5, 6].)

The Δ -category \mathcal{S} is symmetric monoidal under disjoint union and

$$\Omega BS \simeq \mathbb{Z} \times B\Gamma_\infty^+$$

[22, Theorem 3.1].

As the inclusion respects the symmetric monoidal structure, we have the following proposition.

PROPOSITION 2.5. *The inclusion of categories $\mathcal{S} \rightarrow \mathcal{T}$ induces a map of infinite loop spaces $BS \rightarrow B\mathcal{T}$.*

Theorem 4.1 states that $\Omega B\mathcal{T} \simeq \mathbb{Z} \times B\text{Aut}_\infty^+$. The map of infinite loop spaces $\Omega BS \rightarrow \Omega B\mathcal{T}$ is then the map announced in Theorem 1.1.

3. A \mathcal{T} -diagram

In this section, we work with simplicial sets and bisimplicial sets – the latter coming from homotopy colimits of functors to the category of simplicial sets. We use the following fact about bisimplicial sets: to any bisimplicial set $X_{\bullet, \bullet}$, one can associate the *diagonal simplicial set* dX_{\bullet} by taking $dX_p = X_{p,p}$. Let $f : X_{\bullet, \bullet} \rightarrow Y_{\bullet, \bullet}$ be a bisimplicial map. If $f_{p, \bullet}$ is a homotopy equivalence on the vertical simplicial sets $X_{p, \bullet}$ and $Y_{p, \bullet}$ for each $p \geq 0$, then the induced map $f : dX_{\bullet} \rightarrow dY_{\bullet}$ is also a homotopy equivalence [8, IV Lem. 2.6].

When we consider a bisimplicial set as simplicial set, we mean its associated diagonal simplicial set.

Given a small Δ -category \mathcal{M} , an \mathcal{M} -diagram is a functor $\mathcal{X} : \mathcal{M}^{\text{op}} \rightarrow \text{Simp}$, from the opposite category of \mathcal{M} to the category of simplicial sets. It is thus a simplicial set $\mathcal{X} = \coprod_i \mathcal{X}(i)$ where i runs over the objects of \mathcal{M} , with a simplicial action

$$\mathcal{M}(i, j) \times \mathcal{X}(j) \longrightarrow \mathcal{X}(i)$$

for all objects i, j of \mathcal{M} . An example of an \mathcal{M} -diagram is given by taking $\mathcal{X}(i) = \mathcal{M}(i, i_0)$ for a fixed object i_0 of \mathcal{M} .

We denote by \mathcal{T}_1 the \mathcal{T} -diagram with $\mathcal{T}_1(k) := \mathcal{T}(k, 1)$ and by \mathcal{T}_{∞} the \mathcal{T} -diagram with

$$\mathcal{T}_{\infty}(k) = \text{Tel}(\mathcal{T}(k, 1) \xrightarrow{\square \mathbb{G}} \mathcal{T}(k, 1) \xrightarrow{\square \mathbb{G}} \dots),$$

with the maps in the telescope defined by composition with $\mathbb{G} \in \mathcal{T}(1, 1)$, and the \mathcal{T} -diagram structure, as for \mathcal{T}_1 , obtained by pre-composition in \mathcal{T} .

Let $A_{\infty, k} := \text{colim}(A_{n, k} \rightarrow A_{n+1, k} \rightarrow \dots)$, where the map $A_{n, k} \rightarrow A_{n+1, k}$ is given by wedging with an S^1 and extending the homotopies via the identity.

THEOREM 3.1. $\mathcal{T}_{\infty}(k) \simeq \mathbb{Z} \times \text{BA}_{\infty, k}$.

Proof. Recall from Corollary 2.3 that $\mathfrak{T}(k, 1) = \mathfrak{T}_D(k, 1) \sqcup \mathfrak{T}_S(k, 1)$, where $\mathfrak{T}_D(k, 1)$ is the full subcategory of $\mathfrak{T}(k, 1)$ generated by objects with a disc as outgoing surface component. Let $\mathcal{T}_D(k, 1) := N_{\bullet} \mathfrak{T}_D(k, 1)$ and $\mathcal{T}_S(k, 1) := N_{\bullet} \mathfrak{T}_S(k, 1)$ denote their nerves.

Let $\mathcal{T}_{D, \infty}(k) \subset \mathcal{T}_{\infty}(k)$ be the restriction of the telescope $\mathcal{T}_{\infty}(k)$ to the space $\mathcal{T}_D(k, 1)$. As $\mathcal{T}_D(k, 1) \simeq \coprod_{n \geq \varepsilon} \text{BA}_{n, k}$ (Corollary 2.3), we have

$$\mathcal{T}_{D, \infty}(k) \simeq \mathbb{Z} \times \text{BA}_{\infty, k}.$$

The theorem then follows from the fact that there is a retraction

$$r : \mathcal{T}_{\infty}(k) \xrightarrow{\simeq} \mathcal{T}_{D, \infty}(k)$$

as the maps defining the telescope take $\mathcal{T}_S(k, 1)$ to $\mathcal{T}_D(k, 1)$:

$$\mathcal{T}_{\infty}(k) = \text{Tel} \left(\begin{array}{ccccccc} \mathcal{T}_S(k, 1) & & \mathcal{T}_S(k, 1) & & \mathcal{T}_S(k, 1) & & \dots \\ \vdots & \searrow \square \mathbb{G} & \vdots & \searrow \square \mathbb{G} & \vdots & \searrow \square \mathbb{G} & \vdots \\ \mathcal{T}_D(k, 1) & \xrightarrow{\square \mathbb{G}} & \mathcal{T}_D(k, 1) & \xrightarrow{\square \mathbb{G}} & \mathcal{T}_D(k, 1) & \longrightarrow & \dots \end{array} \right)$$

Explicitly, r is a bisimplicial map defined vertically by $r_p : (\mathcal{T}_{\infty}(k))_p = \coprod_{n_0 \leq \dots \leq n_p \in \mathbb{N}} (\mathcal{T}_S(k, 1) \sqcup \mathcal{T}_D(k, 1)) \rightarrow (\mathcal{T}_{D, \infty}(k))_p = \coprod_{n_0 \leq \dots \leq n_p \in \mathbb{N}} \mathcal{T}_D(k, 1)$ which

maps $\mathcal{T}_S(k, 1) \times (n_0, \dots, n_0, n_1, \dots, n_{p-i})$ with $n_0 < n_1$ to $\mathcal{T}_D(k, 1) \times (n_0 + 1, \dots, n_0 + 1, n_1, \dots, n_{p-i})$ using the map $\square_{\mathbb{G}}$ and maps $\mathcal{T}_D(k, 1) \times (n_0, \dots, n_p)$ to itself via the identity. \square

For an \mathcal{M} -diagram \mathcal{X} , we can consider the Borel construction $E_{\mathcal{M}}\mathcal{X}$ of the action, that is the homotopy colimit of the functor \mathcal{X} . Thus $E_{\mathcal{M}}\mathcal{X}$ is the bisimplicial set whose simplicial set of p -simplices is given by

$$(E_{\mathcal{M}}\mathcal{X})_p := \coprod_{i_0, \dots, i_p \in \text{Ob}(\mathcal{M})} \mathcal{M}(i_0, i_1) \times \dots \times \mathcal{M}(i_{p-1}, i_p) \times \mathcal{X}(i_p)$$

with boundary maps d_0 by dropping, d_1, \dots, d_{p-1} by composition in \mathcal{M} and d_p using the \mathcal{M} -diagram structure.

LEMMA 3.2. $E_{\mathcal{T}}\mathcal{T}_{\infty}$ is contractible.

Proof. By [22, Lemma 3.3], $E_{\mathcal{T}}\mathcal{T}_1$ is contractible, where \mathcal{T}_1 is the \mathcal{T} -diagram with $\mathcal{T}_1(n) = \mathcal{T}(n, 1)$. Then $E_{\mathcal{T}}\mathcal{T}_{\infty} \simeq \text{Tel}(E_{\mathcal{T}}\mathcal{T}_1 \rightarrow E_{\mathcal{T}}\mathcal{T}_1 \rightarrow \dots)$ is a telescope of contractible spaces and thus is itself contractible. \square

4. Homotopy type of \mathfrak{T}

We use the \mathcal{T} -diagram \mathcal{T}_{∞} and the homological stability of the automorphisms of free groups with boundary $A_{n,k}$ [13] to prove the following theorem.

THEOREM 4.1. $\Omega B\mathcal{T} \simeq \mathbb{Z} \times \text{BAut}_{\infty}^+$.

LEMMA 4.2. The vertices of the simplicial sets of morphisms in \mathcal{T} act by homology isomorphisms on \mathcal{T}_{∞} .

Proof. The vertices of $\mathcal{T}(n, m)$ are the objects of the category $\mathfrak{T}(n, m)$, so they are 2-dimensional CW-complexes built out of the pieces $\mathbb{P}, \mathbb{T}, \mathbb{D}$ and \mathbb{G} by gluing and disjoint union. It is enough to show that each of these building blocks acts by homology isomorphisms. An element of $\mathcal{T}_{\infty}(k) \simeq \mathbb{Z} \times \text{BA}_{\infty,k}$ is a colimit of morphisms from k to 1 and \mathcal{T} acts by precomposing, that is by gluing on the k incoming boundary circles. Acting one piece at a time means precomposing with that piece disjoint union with $k - 1$ circles. We show that the corresponding operations on the groups $A_{n,k}$ induce homology isomorphisms stably.

By [13], we know that the maps

$$A_{n,k} \xrightarrow{G_{1,0}} A_{n+1,k}$$

$$A_{n,k} \xrightarrow{G_{0,1}} A_{n,k+1}$$

obtained by wedging a circle (that is the graph $G_{1,0}$) and a boundary circle (that is the graph $G_{0,1}$), induce homology isomorphisms on the stable groups:

$$(G_{1,0})_* : H_*(A_{\infty,k}) \xrightarrow{\cong} H_*(A_{\infty,k}) \quad \text{and} \quad (G_{0,1})_* : H_*(A_{\infty,k}) \xrightarrow{\cong} H_*(A_{\infty,k+1}).$$

This implies that gluing a disc \mathbb{D} also induces a homology isomorphism stably since the composition

$$A_{n,k-1} \xrightarrow{G_{0,1}} A_{n,k} \xrightarrow{\mathbb{D}} A_{n,k-1},$$

gluing \mathbb{D} on the boundary circle of the added $G_{0,1}$, is the identity.

Thus gluing a pair of pants \mathbb{P} induces a homology isomorphism stably as the composition with gluing discs on the incoming boundaries of \mathbb{P} is the same as gluing a disc \mathbb{D} where \mathbb{P} was glued.

Gluing a torus \mathbb{T} also induces a homology isomorphism stably as the composition

$$A_{n,k-1} \xrightarrow{G_{0,1}} A_{n,k} \xrightarrow{\mathbb{T}} A_{n+2,k}$$

gluing \mathbb{T} on the boundary circle of $G_{0,1}$, is the same as wedging the graph $G_{2,1}$. Indeed, the two maps add two extra circles and a boundary circle and prolong the homotopy equivalences via the identity on the added pieces. Similarly, gluing \mathbb{G} induces a homology isomorphism stably as the composition with wedging $G_{0,1}$ is the same as wedging $G_{1,1}$. \square

Note that the three maps $A_{n,k} \rightarrow A_{n+2,k}$ induced by gluing \mathbb{T} , gluing $\mathbb{G}\square\mathbb{G}$ or wedging $G_{2,0}$, are all different. However, by freeing a boundary circle (which induces a homology isomorphism stably) one can show that the compositions $F \circ \mathbb{T}$ and $F \circ G_{0,2} : A_{n,k} \rightarrow A_{n+2,k} \rightarrow A_{n+3,k-1}$, gluing a torus or wedging two circles and then freeing the appropriate boundary circle, are conjugate, and hence induce the same map on homology. This is another way to see that gluing \mathbb{T} induces a homology isomorphism stably.

Proof of Theorem 4.1. Consider the following square.

$$\begin{array}{ccc} \mathcal{T}_\infty(k) & \longrightarrow & E_{\mathcal{T}}\mathcal{T}_\infty \\ \downarrow & & \downarrow p \\ k & \longrightarrow & BT \end{array}$$

By the generalized group completion theorem [22, Theorem 3.2] (see also [19, 20]) and using Lemma 4.2, this square is homology cartesian, which means that the fibre of p at any vertex is homology equivalent to the homotopy fibre. As $E_{\mathcal{T}}\mathcal{T}_\infty$ is contractible (Lemma 3.2), the homotopy fibre is ΩBT . Considering the fibre of p at 0, we thus have a homology equivalence $\mathcal{T}_\infty(0) \rightarrow \Omega BT$, and hence again a homology equivalence after plus-construction with respect to any perfect subgroup of $\pi_1\mathcal{T}_\infty(0)$. Now $\mathcal{T}_\infty(0) \simeq \mathbb{Z} \times \text{BAut}_\infty$ by Theorem 3.1 and the commutator subgroup $[\pi_1\mathcal{T}_\infty(0), \pi_1\mathcal{T}_\infty(0)]$ is perfect [18, Remark 2]. Thus we can plus-construct with respect to this subgroup and we get a homotopy equivalence by Whitehead’s theorem for simple spaces [3, Example 4.2]. \square

5. Two infinite loop space structures on $\mathbb{Z} \times \text{BAut}_\infty^+$

Consider the category $\mathfrak{T}_D(0,1)$. Its classifying space is homotopy equivalent to $\coprod_{n \geq 0} \text{BAut}(F_n)$ by Corollary 2.3. We claim that the pair of pants defines a symmetric monoidal structure (up to homotopy) on $\mathfrak{T}_D(0,1)$ equivalent to the symmetric monoidal structure induced by wedging circles on $\coprod_{n \geq 0} \text{Aut}(F_n)$,

thought of as a category with objects $\{\vee_n S^1 \mid n \in \mathbb{N}\}$ and morphism sets $\pi_0 \text{Htpy}_*(\vee_n S^1, \vee_m S^1)$. To be able to remove the ‘up to homotopy’ in the above claim, but also for the comparison to the infinite loop space structure of ΩBT , we need to work with a quotient of the 2-category \mathfrak{T} .

To make the pair of pants multiplication unital, we need to collapse the 1-morphisms $\mathbb{D}\square_i\mathbb{P}$ to the circle, where $i = 1, 2$ represents the two possible gluings of \mathbb{D} and \mathbb{P} . To make it associative, we need to identify $\mathbb{P}\square_1\mathbb{P}$ and $\mathbb{P}\square_2\mathbb{P}$. We define a 2-category \mathfrak{T}^r whose objects are the natural numbers as in \mathfrak{T} , and whose 1-morphisms are the 1-morphisms of \mathfrak{T} with no occurrence of the sequences $\mathbb{D}\square_i\mathbb{P}$ for $i = 1, 2$, or $\mathbb{P}\square_2\mathbb{P}$. The 2-morphisms of \mathfrak{T}^r are as in \mathfrak{T} except that the circle is thought of as a small cylinder, having thus automorphism group \mathbb{Z} . As shown in [25, Section 3.1.1], for each object T of $\mathfrak{T}(n, m)$ there exists a canonical quotient object T^r of $\mathfrak{T}^r(n, m)$ and a homotopy equivalence $T \rightarrow T^r$ which can be used to define composition in \mathfrak{T}^r . (The argument in [25] extends immediately from \mathcal{S} to \mathcal{T} as the quotient construction only affects the pieces \mathbb{P}, \mathbb{T} and \mathbb{D} .) We have $\Omega BT^r \simeq \mathbb{Z} \times \text{BAut}_\infty^+$ by the same argument as for \mathcal{T} . We chose to work with \mathfrak{T} rather than \mathfrak{T}^r in the first part of the paper to stay as simple as possible.

The pair of pants induces a monoidal structure on $\mathfrak{T}_D^r(0, 1)$, where $\mathfrak{T}_D^r(0, 1)$ is defined in an analogous way to $\mathfrak{T}_D(0, 1)$. (Note that the quotient construction only affects the disc \mathbb{D} and not the disc component in \mathbb{G} .) The twist on the pair of pants defines a symmetry as it squares to a Dehn twist along the outgoing boundary of the pair of pants, which is homotopically trivial once the incoming boundaries of the pair of pants are closed by discs.

THEOREM 5.1. *The equivalence $\Omega BT \simeq \mathbb{Z} \times \text{BAut}_\infty^+$ is an equivalence of infinite loop spaces, where the left infinite loop structure is induced by disjoint union in \mathcal{T} and the right one by wedging circles in $\coprod_{n \in \mathbb{N}} \text{Aut}(F_n)$.*

Proof. The proof of this theorem is totally analogous to the proof of the main result in [25], which says that the infinite loop space structure on $\mathbb{Z} \times \text{B}\Gamma_\infty^+$ coming from $\Omega B\mathcal{S}$ is equivalent to the one coming from the \mathcal{M} -algebra structure of $\mathcal{S}(0, 1) \simeq \coprod_{g \geq 0} \text{B}\Gamma_{g,1}$, where \mathcal{M} is the mapping class group operad defined in [24]. We will only sketch the main steps for the case of interest here.

We have an equivalence of infinite loop spaces

$$\Omega BT \xrightarrow{\simeq} \Omega BT^r,$$

where \mathcal{T}^r denotes the Δ -category associated to \mathfrak{T}^r , and we have an equivalence of symmetric monoidal categories

$$\mathfrak{T}_D^r(0, 1) \xrightarrow{\simeq} \coprod_{n \in \mathbb{N}} \text{Aut}(F_n).$$

We give here an equivalence between the spectra of deloops of $\Omega B(\mathcal{T}_D^r(0, 1))$ and of ΩBT^r .

Let Σ be the standard E_∞ operad with $\Sigma(k) = E\Sigma_k$ for Σ_k the symmetric group. (This operad is denoted Γ^+ in [1, 2] and Γ in [25].) As $\mathfrak{T}_D^r(0, 1)$ and \mathcal{T}^r are symmetric monoidal categories, the operad Σ acts on their classifying spaces $\mathcal{T}_D^r(0, 1)$ and $B\mathcal{T}^r$.

We want to relate ΩBT^r and $\mathcal{T}_D^r(0, 1)$, which is a subspace of $\mathcal{T}^r(0, 1)$. To a morphism from 0 to 1 in \mathcal{T}^r corresponds a 1-simplex with boundary points 0 and




FIGURE 4. Homotopy.

1 in BT^r , by definition of the nerve. There is thus a natural map

$$\phi : \mathcal{T}_D^r(0, 1) \rightarrow \Omega BT^r$$

taking the path from 0 to 1 in BT^r defined by the element of $\mathcal{T}_D^r(0, 1)$ and going back to 0 along the path defined by the disc, also thought of as a morphism from 0 to 1. The crucial observation is that this map respects the multiplication up to homotopy, as show in Figure 4. The figure shows the loop obtained by multiplying $\mathbb{D} \square \mathbb{T} \square \mathbb{G}$ and $\mathbb{D} \square \mathbb{G}$ in $\mathcal{T}_D^r(0, 1)$ and then mapping to ΩBT^r , producing a loop following $(\mathbb{D} \square \mathbb{T} \square \mathbb{G} \sqcup \mathbb{D} \square \mathbb{G}) \square \mathbb{P}$ from 0 to 1, and then following the disc-morphism back to 0. The other loop, going from 0 to 2 and back to 0, is the one obtained by mapping first $\mathbb{D} \square \mathbb{T} \square \mathbb{G}$ and $\mathbb{D} \square \mathbb{G}$ to ΩBT^r and then multiplying, that is taking loop on disjoint union. The two loops are homotopic in BT^r because the operad Σ acts on $\mathcal{T}_D^r(0, 1)$ by taking disjoint union and composing with \mathbb{P} as a morphism in \mathcal{T}^r . This means that the top triangle in the figure commutes in \mathcal{T}^r , and hence defines a 2-simplex in BT^r . On the other hand, we have defined composition in \mathcal{T}^r so that the bottom triangle also commutes. (This is a place where we need the disc to be a strict unit.)

This argument extends to give all necessary higher homotopies, parametrized by the elements of Σ . (Considering higher multiplications will give higher simplices in BT^r .)

More concretely, we consider Barratt and Eccles' model for the spectra associated to the action of Σ on $\mathcal{T}_D^r(0, 1)$ and on BT^r . These are realization of simplicial spaces, where the i th space of the spectrum has q -simplices given by $\mathcal{G}\Sigma(S^i \wedge \Sigma^q(\mathcal{T}_D^r(0, 1)))$ and $\mathcal{G}\Sigma(S^i \wedge \Sigma^q(BT^r))$ respectively, where \mathcal{G} is the group completion functor on free simplicial monoids, Σ denotes now the monad associated to the operad Σ , and Σ^q means iterating the monad q times. The simplicial structure is given by the monad multiplication for d_0, \dots, d_{q-1} (using the assembly map $S^1 \wedge \Sigma(X) \rightarrow \Sigma(S^1 \wedge X)$

for d_0) and by the Σ -algebra structure of $\mathcal{T}_D^r(0, 1)$ and BT^r for the last boundary map. The spectrum structure comes from the equivalence

$$\mathcal{G}\Sigma(X) \xrightarrow{\simeq} \Omega\mathcal{G}\Sigma(S^1 \wedge X)$$

[1, 2].

The map ϕ defined above gives a map on the level of q -simplices from the i th deloop to the $(i - 1)$ th deloop.

$$\begin{array}{ccc} \mathcal{G}\Sigma(S^i \wedge \Sigma^q(\mathcal{T}_D^r(0, 1))) & \xrightarrow{f_q^i} & \mathcal{G}\Sigma(S^{i-1} \wedge \Sigma^q(BT^r)) \\ \downarrow & \nearrow \phi & \\ \mathcal{G}\Sigma(S^{i-1} \wedge \Sigma^q(S^1 \wedge \mathcal{T}_D^r(0, 1))) & & \end{array}$$

The maps $\{f_q^i\}_{q \geq 0}$ do not quite form a simplicial map as ϕ respects the algebra structure only up to homotopy, as explained above. However, as in the case of the mapping class groups, one can use the explicit homotopies sketched above to rectify f_*^i to a simplicial map, while respecting the spectrum structure. The rectified map gives the equivalence of spectra. \square

Proposition 2.5, Theorem 4.1 and Theorem 5.1 combine to prove Theorem 1.1.

6. Punctured surfaces

We have constructed a 2-category \mathfrak{T} which is adequate for proving Theorem 1.1. There are of course many possible versions of \mathfrak{T} . M. Weiss suggested the following modification of \mathfrak{T} , which has the advantage of being a little more natural to construct, but the disadvantage of losing basepoints, thus making the comparison to the original infinite loop space structure on $\mathbb{Z} \times \text{BAut}_\infty^+$ more difficult.

Let \mathfrak{D} be the 2-category obtained from \mathfrak{T} by replacing the 1-morphism \mathbb{G} by a punctured cylinder $\mathbb{C} = (S^1 \times I) \setminus \{*\}$, with one incoming and one outgoing boundary circle. The 1-morphisms of \mathfrak{D} are (possibly punctured) cobordisms built out of $\mathbb{P}, \mathbb{T}, \mathbb{D}$ and \mathbb{C} , and the 2-morphisms are homotopy equivalences which fix the boundary circles, but not the punctures.

As soon as a surface is punctured, it is homotopy equivalent to a basepoint-free graph $G_{n,k}$, relative to the boundary circles. Let $A_{n,k}^0 = \pi_0 \text{Htpy}(G_{n,k}; \partial)$ denote the group of components of the space of homotopy equivalences of $G_{n,k}$ which fix its k boundary circles. Let $\mathfrak{D}_p(k, 1)$ denote the subcategory of $\mathfrak{D}(k, 1)$ generated by the punctured surfaces. We have

$$\mathcal{D}_p(k, 1) := \text{B}\mathfrak{D}_p(k, 1) \simeq \coprod_{n \geq 0} \text{BA}_{n,k+1}^0.$$

As the groups $A_{n,k}^0$ have the same stable homology as $A_{n,k}$ [13], one can run through Sections 2–4 above replacing \mathfrak{T} by \mathfrak{D} and \mathbb{G} by \mathbb{C} , and show that $\Omega \text{BD} \simeq \mathbb{Z} \times \text{BAut}_\infty^+$. Only Section 5 does not have a straightforward extension.

Let \mathcal{M} denote the mapping class group operad of [24]. One can define \mathcal{M} in terms of the cobordism category by taking $\mathcal{M}(k) = \mathcal{S}(k, 1)$ and the operad composition induced by composition in \mathcal{S} . It is shown in [24] that \mathcal{M} -algebras are infinite loop spaces after group completion. The pair of pants multiplication does not define a symmetric monoidal structure on $\mathcal{D}_p(0, 1)$, but it extends to an action of \mathcal{M} . One

can adapt the proof of Theorem 5.1 (or rather the proof of the main theorem of [25]) to show the equivalence between the infinite loop space structure of ΩBD , induced by disjoint union on \mathcal{D} , and that of $\Omega BD_p(0, 1)$, induced by the \mathcal{M} -algebra structure of $\mathcal{D}_p(0, 1)$. However, the only method I can see for comparing the infinite loop space structure of $\Omega BD_p(0, 1)$ and the ‘usual’ structure of $\mathbb{Z} \times \text{BAut}_\infty^+$ is going through ΩBD and ΩBT via a middle category with both \mathbb{G} and \mathbb{C} as 1-morphisms.

$$\begin{array}{ccccc}
 \Omega BD_p(0, 1) & \longrightarrow & \Omega BD & & \Omega B \coprod_{n \in \mathbb{N}} \text{BAut}(F_n) \\
 & & \downarrow & & \uparrow \\
 & & \Omega B(\mathcal{D} \cup \mathcal{T}) & \longleftarrow & \Omega BT \longleftarrow \Omega BT_D(0, 1)
 \end{array}$$

Note that $\mathcal{D}_p(0, 1)$ is a sub- \mathcal{M} -algebra of $\mathcal{D}(0, 1)$, and one can show that they have the same group completion, namely $\mathbb{Z} \times \text{BAut}_\infty^+$. We thus have a map of \mathcal{M} -algebras

$$S(0, 1) \longrightarrow \mathcal{D}(0, 1)$$

which after group completion gives another model for the infinite loop map

$$\mathbb{Z} \times B\Gamma_\infty^+ \longrightarrow \mathbb{Z} \times \text{BAut}_\infty^+.$$

Acknowledgements. I would like to thank the Aarhus group, and in particular Marcel Böksted, Søren Galatius and Ib Madsen for many helpful conversations. I would also like to thank Allen Hatcher and Craig Jensen for their collaboration on the related papers [13] and [15], and Benson Farb and Ulrike Tillmann for conversations at the early stages of this paper. Finally, I would like to thank the referee for suggesting a nice improvement to the paper.

References

1. M. G. BARRATT and P. J. ECCLES, ‘ Γ^+ -structures. I: A free group functor for stable homotopy theory’, *Topology* 13 (1974) 25–45.
2. M. G. BARRATT and P. J. ECCLES, ‘ Γ^+ -structures. II: A recognition principle for infinite loop spaces’, *Topology* 13 (1974) 113–126.
3. E. DROR, ‘A generalization of the Whitehead theorem’, *Symposium on Algebraic Topology (Battelle Seattle Research Center, Seattle, WA, 1971)*, Lecture Notes in Mathematics 249 (Springer, Berlin, 1971) 13–22.
4. W. DWYER, M. WEISS and B. WILLIAMS, ‘A parametrized index theorem for the algebraic K -theory Euler class’, *Acta Math.* 190 (2003) 1–104.
5. C. J. EARLE and J. EELLS, ‘A fibre bundle description of Teichmüller theory’, *J. Differential Geom.* 3 (1969) 19–43.
6. C. J. EARLE and A. SCHATZ, ‘Teichmüller theory for surfaces with boundary’, *J. Differential Geom.* 4 (1970) 169–185.
7. S. GALATIUS, ‘Mod p homology of the stable mapping class group’, *Topology* 43 (2004) 1105–1132.
8. P. G. GOERSS and J. F. JARDINE, *Simplicial homotopy theory*, Progress in Mathematics 174 (Birkhäuser, Basel, 1999).
9. J. L. HARER, ‘Stability of the homology of the mapping class groups of orientable surfaces’, *Ann. of Math.* (2) 121 (1985) 215–249.
10. A. HATCHER, *Algebraic topology* (Cambridge University Press, Cambridge, 2002).
11. A. HATCHER and K. VOGTMANN, ‘Cerf theory for graphs’, *J. London Math. Soc.* (2) 58 (1998) 633–655.
12. A. HATCHER and K. VOGTMANN, ‘Rational homology of $\text{Aut}(F_n)$ ’, *Math. Res. Lett.* 5 (1998) 759–780.
13. A. HATCHER and N. WAHL, ‘Stabilization for the automorphisms of free groups with boundaries’, arXiv:math.GT/0406277.
14. N. V. IVANOV, ‘Stabilization of the homology of Teichmüller modular groups’, *Algebra i Analiz* 1 (1989) 110–126.

15. C. A. JENSEN and N. WAHL, ‘Automorphisms of free groups with boundaries’, *Algebr. Geom. Topol.* 4 (2004) 543–569.
16. I. MADSEN and M. S. WEISS, ‘The stable moduli space of Riemann surfaces: Mumford’s conjecture’, arXiv:math.AT/0212321.
17. J. P. MAY, ‘ E_∞ spaces, group completions, and permutative categories’, *New developments in topology (Proceedings of Symposium on Algebraic Topology, Oxford, 1972)*, London Mathematical Society Lecture Note Series 11 (Cambridge University Press, London, 1974) 61–93.
18. D. MCDUFF and G. SEGAL, ‘Homology fibrations and the “group-completion” theorem’, *Invent. Math.* 31 (1975/76) 279–284.
19. I. MOERDIJK, ‘Bisimplicial sets and the group-completion theorem’, *Algebraic K-theory: connections with geometry and topology (Lake Louise, AB, 1987)*, NATO Advanced Science Institutes Series C: Mathematical and Physical Sciences 279 (Kluwer, Dordrecht, 1989) 225–240.
20. W. PITSCH and J. SCHERER, ‘Homology fibrations and “group-completion” revisited’, *Homology Homotopy Appl.* 6 (2004) 153–166 (electronic).
21. G. SEGAL, ‘Categories and cohomology theories’, *Topology* 13 (1974) 293–312.
22. U. TILLMANN, ‘On the homotopy of the stable mapping class group’, *Invent. Math.* 130 (1997) 257–275.
23. U. TILLMANN, ‘A splitting for the stable mapping class group’, *Math. Proc. Cambridge Philos. Soc.* 127 (1999) 55–65.
24. U. TILLMANN, ‘Higher genus surface operad detects infinite loop spaces’, *Math. Ann.* 317 (2000) 613–628.
25. N. WAHL, ‘Infinite loop space structure(s) on the stable mapping class group’, *Topology* 43 (2004) 343–368.

Nathalie Wahl
Department of Mathematics
University of Aarhus
Ny Munkegade 116
8000 Aarhus
Denmark
wahl@imf.au.dk