

# Algebra I/Introduction to Algebra V

Due: Please upload solutions to TACT by Tuesday, June 20, 2023.

**Problem 1.** Let  $G = \Sigma_4$  be the symmetric group on 4 letters, and recall that  $G$  has 5 conjugacy classes of elements with representatives  $x_1 = e$ ,  $x_2 = (12)$ ,  $x_3 = (12)(34)$ ,  $x_4 = (123)$ , and  $x_5 = (1234)$ . The number  $\ell_i = [G : G_{x_i}]$  of elements in the conjugacy class containing  $x_i$  is  $\ell_1 = 1$ ,  $\ell_2 = 6$ ,  $\ell_3 = 3$ ,  $\ell_4 = 8$ , and  $\ell_5 = 6$ . In Lecture 7, we defined 5 pairwise non-isomorphic irreducible finite dimensional complex representations of  $G$ , namely, the trivial 1-dimensional representation  $\pi_1$ , the 1-dimensional sign representation  $\pi_2$ , the 3-dimensional standard representation  $\pi_3$ , the 3-dimensional representation  $\pi_4 = \pi_2 \otimes \pi_3$ , and a 2-dimensional representation  $\pi_5$  obtained from the standard representation  $\bar{\pi}_3$  of  $\bar{G} = \Sigma_3$ .<sup>1</sup> Let  $\chi_i$  be the character of  $\pi_i$ . By using the definition of  $\pi_1$ ,  $\pi_2$ , and  $\pi_3$  it is (rather) easy to find the values of  $\chi_1$ ,  $\chi_2$ , and  $\chi_3$  on the  $x_1, \dots, x_5$ :

$x_i$	$e$	$(12)$	$(12)(34)$	$(123)$	$(1234)$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	-1	1	1	-1
$\chi_3$	3	1	-1	0	-1
$\chi_4$	3	?	?	?	?
$\chi_5$	2	?	?	?	?
$\ell_i$	1	6	3	8	6

Please answer the following questions:

- (a) Find the values of  $\chi_4$  on  $x_1, \dots, x_5$ .

[Hint: Use that  $\chi_{\pi \otimes \tau} = \chi_\pi \cdot \chi_\tau$ .]

- (b) Find the values of  $\chi_5$  on  $x_1, \dots, x_5$ .

[Hint: Use that  $(\chi_1, \chi_2, \chi_3, \chi_4, \chi_5)$  is a basis of  $Z(\mathbb{C}[G])$  and orthonormal with respect to the Schur inner product to produce a system of 4 linear equations in the 4 variables  $\chi_5(x_2)$ ,  $\chi_5(x_3)$ ,  $\chi_5(x_4)$ , and  $\chi_5(x_5)$  and solve it.]

- (c) Find the non-negative integers  $m_1, \dots, m_5$  such that

$$\pi_3 \otimes \pi_3 \simeq \pi_1^{m_1} \oplus \pi_2^{m_2} \oplus \pi_3^{m_3} \oplus \pi_4^{m_4} \oplus \pi_5^{m_5}.$$

Here  $\pi^m$  denotes the direct sum of  $m$  copies of  $\pi$ .

[Hint: Use that  $(\chi_1, \chi_2, \chi_3, \chi_4, \chi_5)$  is a basis of  $Z(\mathbb{C}[G])$  that is orthonormal with respect to the Schur hermitian inner product.]

---

<sup>1</sup> In the language of Lecture 9, we defined a group homomorphism  $p: G = \Sigma_4 \rightarrow \bar{G} = \Sigma_3$  and defined  $\pi_5 = p^*(\bar{\pi}_3)$  to be the representation of  $G$  obtained from the (2-dimensional) standard representation  $\bar{\pi}_3$  of  $\bar{G}$  by restriction along  $p$ .